
MatriMix

New 3D Culture Substrate

MatriMix for 3D cell culture

MatriMix is a new 3D culture substrate consisting of collagen, laminin E8 fragments (LM E8) and hyaluronan. The type, combinations, and concentrations of each ingredient can be customized to provide a microenvironment that is suitable for various types of cells. MatriMix is composed of 3 solutions (A; DMEM/LM E8-crosslinked hyaluronan, B; sodium bicarbonate, C; collagen), which are mixed just before incubation. The mixture, which is a solution in cold storage, gels by warming to 37 °C. MatriMix can be used not only for "in/on gel" cell culture, but also for cell transplantation into mice.



The laminin C-terminal E8 fragment, which is about 1/5 of the full-length laminin molecule, is recombinantly expressed. Laminin-511E8 has a strong interaction with cellular integrin $\alpha 6\beta 1$ and induces cell motility.





AMS Biotechnology (Europe) Ltd UK & Rest of the World

184 Park Drive, Milton Park Abington OX14 4SE. T: +44 (0) 1235 828 200 F: +44 (0) 1235 820 482

AMSBIO LLC

USA & Canada 1035 Cambridge Street, Cambridge, MA 02141. T: +1 (617) 945-5033 or T: +1 (800) 987-0985

F: +1 (617) 945-8218

AMSBIO Europe BV Europe

Berenkoog 41, 1822 BH Alkmaar, Netherlands T: +31 (0) 72 8080244 F: +31 (0) 72 8080142

•

AMS Biotechnology (Europe) Ltd Switzerland Via Lisano 3,


VIA LISANO 3, (CP.683) CH-6900 T: +41 (0) 91 604 55 22 F: +41 (0) 91 605 17 85



Bockenheimer Landstr. 17/19 60325 Frankfurt/Main T: +49 (0) 69 779099 F: +49 (0) 69 13376880


Induction of organoid formation by cancer cells and early embryo derived cells

Drug screening




- A new, a more "in vivo mimetic" in vitro model that solves various problems with the EHS tumor extracts.
- Excellent for cultured cells that are difficult to grow and organize in the EHS tumor extracts.
- Drug screening can be performed in vitro without transplantation into a mouse model.
- Drugs can be screened for highly malignant cancers with metastatic potential.

Organoid research



Developed especially for scientists not satisfied with current 3D substrates

Effects of MatriMix gels on colon cancer patients derived spheroid cultures



ZEB-1 (green), E-cad (red), DAPI (blue)

In MatriMix gels, cells derived from patients with Stage 4 metastatic cancer showed a cell population positive for the metastatic potential marker ZEB-1. In contrast, cells derived from Stage 3 patients with non-metastasis cancer demonstrated a suppression of ZEB-1 expression.



MatriMix gel can be used for the transplantation of cancer spheroids into nude mice similar to EHS tumor extracts.

For more information and for other applications, please contact us at: info@amsbio.com