

Anti Human Interleukin-4 (IL-4) Azide Free Monoclonal Mouse Antibody

Code: 211-44-534C **Lot no.:** 1288-1009P
Clone: 8F12 **Exp.:** 1 year from date of dispatch
Isotype: Mouse IgG1
Quantity: 500µg (1 mg/ml) antibody buffered in phosphate buffered saline, pH 8.0 without Sodium Azide.

Purification: Protein A affinity chromatography.

Specificity: This antibody recognizes both recombinant and natural forms of human IL-4¹.

Applications: Flow cytometry: Intracellular staining of IL-4 producing cells with clone 8F12 was demonstrated by flow cytometric analysis. Mitogen stimulated purified T cells treated with saponin and monensin yielded a strong positive signal with clone 8F12 in flow cytometry². Allergen specific T cell clones derived from patients with atopic dermatitis showed a high degree of correlation between IL-4 levels in supernatants by ELISA and intracellular flow cytometric analysis. A correlation was observed with respect to the pattern of cytokine (Th0, Th1 or Th2) production^{1,2}.

Immunocytochemistry (ICC): Intracellular staining of human IL-4 with clone 8F12 was demonstrated with the use of transfected CHO cells expressing IL-4^{3,4}. Clone 8F12 was also used for intracellular staining of mitogen stimulated purified human T cells⁸ and gradient purified human basophils stimulated with IL-3⁴.

ELISA: Biotinylated anti human IL-4 antibody (clone 8F12; cat# 211-44-234C) can be used as the detecting antibody in a two-site sandwich ELISA for measuring human IL-4 levels. The purified form of anti human IL-4 may be used as the capture reagent (clone 3H4; Cat # 211-44-134B)^{1,4,5,6}, with recombinant human IL-4 (Cat # 111-40-134) as the standard. Optimal concentration for the biotin-conjugated detecting monoclonal (clone 8F12) was 2µg/ml. This ELISA measured human IL-4 with a detection limit of 5 pg/ml and range up to 3 ng/ml.

Neutralization: Clone 8F12 was demonstrated to neutralize the biological activity of human IL-4 by inhibition of IL-4 stimulated T cell proliferation⁷. The inhibition of 50% maximal T cell proliferation (IC₅₀) was equal to 0.3 nM of clone 8F12. We recommend the azide free format of clone 8F12 for this application (cat # 211-44-534C).

Western Blotting: Clone 8F12 was found to react equally well with native and denatured amounts of human IL-4 (smaller than 10 ng), in dot blot and Western blot analysis⁷.

References:

1. Jung T. et al. (1995). *Eur. J. Immunol.* 25: 2413-2416.
2. Jung T. et al. (1993). *J. Immunol. Meth.* 159: 197-207.
3. Mueller R. et al. (1994). *Cell Biology International.* 18(1): 55-61.
4. Mueller, R. et al. (1994). *Eur. J. Immunol.* 24: 2935-2940.
5. Brunner, T. et al. (1993). *J. Exp. Med.* 177: 605.
6. Kilchherr, E. et al. (1993). *Cell. Immunol.* 151:241.
7. Reusch, P. et al. (1994). *Eur. J. Biochem.* 222: 491-499.
8. Andersson, U. et al. (1990). *Eur. J. Immunol.* 20: 1591-1596.

Storage: Since antibody solution does not contain sodium azide or other preservatives, we recommend to aliquot solution into small volumes under sterile conditions and store at -20°C.

For Research Use Only. Not For Diagnostic or Therapeutic Use.

Conditions: The information disclosed herein is not to be considered as a recommendation to use the above product in violation of any patents. ImmunoKontakt will not be held responsible for patent infringement or other violations that may occur with the use of our products.

Rev. Sep-12

UK & REST OF WORLD

184 MILTON PARK, ABINGDON
OX14 4SE, OXON, UK
TEL: +44 (0) 1235 828 200
FAX: +44 (0) 1235 820 482

UNITED STATES

23591 EL TORO RD, SUITE #180
LAKE FOREST, CA 92630
TEL: + 1 800 987 0985
FAX: + 1 949 265 7703

SWITZERLAND

CENTRO NORD-SUD 2E
CH-6934 BIOGGIO-LUGANO
TEL: +41 (0) 91 604 55 22
FAX: +41 (0) 91 605 17 85

DEUTSCHLAND

BOCKENHEIMER LANDSTR. 17/19
60325 FRANKFURT/MAIN
TEL: +49 (0) 69 779099
FAX: +49 (0) 69 13376880

