

# **Product Data Sheet**

# HT29 A-FLX™ FFPE Cell Pellet

# GENERAL INFORMATION

Product Name: HT29 A-FLX™ FFPE Cell Pellet

Reference Number: 3050-0210 Block

3050-0220 Slide (5μm)

3050-0230 FFPE scroll (20µm)

Date of Manufacturing: See product label
Lot Number: See product label
Intended Use: For research use only

#### **DESCRIPTION**

Cell Line: HT29

Tissue of Origin: Colon or rectum

Culturing Condition: DMEM supplemented with 10% FBS at 37°C with

5% CO<sub>2</sub>

Fixation Condition: 10% neutral buffered formalin (NBF) for 24 hours

at 24-27°C

Product Format:

Block: Paraffin embedded block. Pellet thickness: ~2mm

Slide: One unstained section mounted on Superfrost™

Plus slide. Section thickness: 5µm

FFPE Scroll: One FFPE section in DNase/RNase free tube.

Section thickness: 20µm

# **SCHEMATICS**




Illustration of an FFPE slide

#### SAFETY AND PRECAUTIONS

This product does not contain hazardous material. Wear appropriate personal productive equipment (PPE) when handling reagents and biological specimens.

## RECOMMENDED PROCEDURES

## Staining using FFPE slides:

- 1. Bake slides at 60°C for 30-60 min.
- 2. Deparaffinize two times in Xylene or Xylene substitute for 5 min each time.
- 3. Rinse two times in 100% ethanol for 1 min each time.
- 4. Air dry slides or rehydrate in ethanol series (95% 2 min, 70% 2 min, 50% 2 min, 1XPBS 2 min).
- 5. Proceed to staining protocol.

#### Biomolecule extraction from FFPE scrolls:

- 1. Add 1ml Xylene or Xylene substitute to each tube containing FFPE scrolls and vortex for 10 sec.
- 2. Centrifuge at full speed for 2 min at room temperature. Remove supernatant without disturbing the pellet.
- 3. Repeat step 1 and 2.
- 4. Add 1ml 100% ethanol and mix by vortexing.
- 5. Centrifuge at full speed for 2 min at room temperature. Remove supernatant without disturbing the pellet.
- 6. Repeat step 4 and 5.
- 7. Carefully remove any residual ethanol in the tube without disturbing the pellet.
- 8. Open the tube and dry at room temperature or up to 37°C for 10min, or until the ethanol has completely evaporated.
- 9. Proceed to extraction protocol.

#### STORAGE AND STABILITY

Storage Condition 2-8°C with desiccation (Recommended)

#### **STABILITY**

Block 3 years
Scroll 1 year
Slide 1 year

AMSBIO | www.amsbio.com | info@amsbio.com



F: +44 (0) 1235 820 482





<sup>\*</sup>Guideline for general applications, such as immunohistochemistry (IHC) or DNA in situ hybridization (ISH). Certain biomolecules may be less stable during storage