amsbio

## HL60 FFPE Cell Pellet

#### GENERAL INFORMATION

Product Name:	HL60 FFPE Cell Pellet	
Reference Number:	3070-0410	Block
	3070-0420	Slide (5µm)
	3070-0430	FFPE scroll (20µm)
Date of Manufacturing:	See product label	
Lot Number:	See product label	
Intended Use:	For research use only	

#### DESCRIPTION

Cell Line:	HL60
Tissue of Origin:	Leukemia
Culturing Condition:	Multiple
Fixation Condition:	10% neutral buffered formalin (NBF) for 24 hours at 24-27°C
Product Format:	
Block:	Paraffin embedded block. Pellet thickness: ~2mm
Slide:	One unstained section mounted on Superfrost™ Plus slide. Section thickness: 5µm
FFPE Scroll:	One FFPE section in DNase/RNase free tube. Section thickness: 20µm

#### SCHEMATICS




Illustration of an FFPE slide

# **Product Data Sheet**

#### SAFETY AND PRECAUTIONS

This product does not contain hazardous material. Wear appropriate personal productive equipment (PPE) when handling reagents and biological specimens.

#### **RECOMMENDED PROCEDURES**

#### Staining using FFPE slides:

- 1. Bake slides at 60°C for 30-60 min.
- 2. Deparaffinize two times in Xylene or Xylene substitute for 5 min each time.
- 3. Rinse two times in 100% ethanol for 1 min each time.
- 4. Air dry slides or rehydrate in ethanol series (95% 2 min, 70% 2 min, 50% 2 min, 1XPBS 2 min).
- 5. Proceed to staining protocol.

#### Biomolecule extraction from FFPE scrolls:

- 1. Add 1ml Xylene or Xylene substitute to each tube containing FFPE scrolls and vortex for 10 sec.
- 2. Centrifuge at full speed for 2 min at room temperature. Remove supernatant without disturbing the pellet.
- 3. Repeat step 1 and 2.
- 4. Add 1ml 100% ethanol and mix by vortexing.
- 5. Centrifuge at full speed for 2 min at room temperature. Remove supernatant without disturbing the pellet.
- 6. Repeat step 4 and 5.
- 7. Carefully remove any residual ethanol in the tube without disturbing the pellet.
- 8. Open the tube and dry at room temperature or up to 37°C for 10min, or until the ethanol has completely evaporated.
- 9. Proceed to extraction protocol.

#### STORAGE AND STABILITY

Storage Condition	2-8°C with desiccation (Recommended)
STABILITY	
Block	3 years
Scroll	1 year
Slide	1 year

\*Guideline for general applications, such as immunohistochemistry (IHC) or DNA in situ hybridization (ISH). Certain biomolecules may be less stable during storage

### AMSBIO| www.amsbio.com | info@amsbio.com

UK & Rest of the World 184 Park Drive, Milton Park Abingdon OX14 4SE. T: +44 (0) 1235 828 200 F: +44 (0) 1235 820 482

North America 1035 Cambridge Street, Cambridge, MA 02141. T: +1 (617) 945-5033 or T: +1 (800) 987-0985 F: +1 (617) 945-8218

Europe Berenkoog 41, 1822 BH Alkmaar, Netherlands T: +31 (0) 72 8080244 F: +31 (0) 72 8080142 Switzerland Via Lisano 3, (CP.683) CH-6900 T: +41 (0) 91 604 55 22 F: +41 (0) 91 605 17 85 Germany Bockenheimer Landstr. 17/19 60325 Frankfurt/Main T: +49 (0) 69 779099 F: +49 (0) 69 13376880