

Data Sheet

TNKS2 Histone Ribosylation Assay Kit (Biotin-labeled NAD⁺) Catalog # 80578

DESCRIPTION: The TNKS2 Histone Ribosylation Assay Kit (Biotin-labeled NAD⁺) is designed to measure Tankyrase 2 (TNKS2) activity for screening and profiling applications. TNKS2 catalyzes the NAD-dependent addition of poly(ADP-ribose) to the substrate proteins. The TNKS2 assay kit comes in a convenient 96-well format, with purified TNKS2 enzyme, histone mixture, and PARP assay buffer for 100 enzyme reactions. The key to the TNKS2 Histone Ribosylation Assay is the biotinylated NAD⁺ substrate. With this kit, only three simple steps are required for TNKS2 reactions. First, histone proteins are coated on a 96-well plate. Next, the biotinylated NAD⁺ substrate is incubated with an assay buffer that contains the TNKS2 enzyme. Finally, the plate is treated with streptavidin-HRP followed by addition of the HRP substrate to produce chemiluminescence that can be measured using a chemiluminescence reader.

COMPONENTS:

Reagent	Amount	Storage
TNKS2 (BPS # 80515)	2 µg	-80 °C
5x histone mixture	1 ml	-80 °C
10x Assay Mixture containing biotinylated substrate	300 µl	-80 °C
10x PARP assay buffer	1 ml	-20 °C
Blocking buffer	25 ml	+4 °C
Streptavidin-HRP	100 µl	+4 °C
HRP chemiluminescent substrate (2 components)	6 ml each	+4 °C
96-well module	1	Room Temperature


MATERIALS AND INSTRUMENTS REQUIRED BUT NOT SUPPLIED:

1x PBS buffer
 PBST buffer (1x PBS, containing 0.05% Tween-20)
 Luminometer or fluorescent microplate reader capable of reading chemiluminescence
 Adjustable micropipettor and sterile tips
 Rotating or rocker platform

APPLICATIONS: Great for studying enzyme kinetics and screening small molecular inhibitors for drug discovery and HTS applications.

STABILITY: Up to 1 year when stored as recommended.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



UK & Rest of World

184 Milton Park, Abingdon
OX14 4SE, Oxon, UK
Tel: +44 (0) 1235 828 200
Fax: +44 (0) 1235 820 482

Switzerland

Centro Nord-Sud 2E
CH-6934 Bioggio-Lugano
Tel: +41 (0) 91 604 55 22
Fax: +41 (0) 91 605 17 85

Deutschland

Bockenheimer Landstr. 17/19
60325 Frankfurt/Main
Tel: +49 (0) 69 779099
Fax: +49 (0) 69 13376880

North America

23591 El Toro Rd, Suite #180
Lake Forest, CA 92630
Tel: +1 800 987 0985
Fax: +1 949 265 7703

amsbio

info@amsbio.com

www.amsbio.com
AMS Biotechnology

REFERENCE:

Brown, J.A., Marala, R.B. *J. Pharmacol. Toxicol. Methods* 2002 **47**:137-41.

Assay Protocol:

All samples and controls should be tested in duplicate.

Coating the plate with the histone mixture:

- 1) Dilute 5x histone mixture 1:5 in PBS.
- 2) Add 50 µl diluted histone mixture to each well and incubate overnight at 4°C.
- 3) Wash the plate 3 times with 200 µl PBST buffer.
- 4) Block the wells by adding 150 µl of Blocking buffer to every well. Incubate for 60 minutes at room temperature.
- 5) Wash the plate 3 times with 200 µl PBST buffer

(Alternatively, the plate can be coated for 90 minutes at 37°C followed by 60 minutes blocking at room temperature. All washing steps should be the same.)

Step 1: Ribosylation reaction

- 1) Prepare the master mixture: N wells x (2.5 µl **10x PARP assay buffer** + 2.5 µl **10x PARP assay mixture** + 20 µl **H₂O**)
- 2) Thaw TNKS2 **enzyme** on ice. Upon first thaw, briefly spin tube containing enzyme to recover full content of the tube. Aliquot TNKS2 **enzyme** into single use aliquots. Store remaining undiluted enzyme in aliquots at -80°C immediately. *Note: TNKS2 enzyme is very sensitive to freeze/thaw cycles. Do not re-use thawed aliquots or diluted enzyme.*
- 3) Dilute TNKS2 **enzyme** in 1X PARP assay buffer at 0.75 – 1 ng/µl (15 - 20 ng/20 µl). Keep diluted enzyme on ice until use. Discard any unused diluted enzyme after use.

Add 25 µl of master mixture to each well designated for the “Positive Control”, “Test Inhibitor”, and “Blank”. For the “Substrate Control”, add 2.5 µl **10x PARP assay buffer** + 22.5 µl **H₂O**.

	Blank	Positive Control	Substrate Control	Test Inhibitor
10X PARP Assay Buffer	2.5 µl	2.5 µl	2.5 µl	2.5 µl
10X assay mixture	2.5 µl	2.5 µl	–	2.5 µl
H ₂ O	20 µl	20 µl	22.5 µl	20 µl
Test Inhibitor	–	–	–	5 µl
Inhibitor buffer (no inhibitor)	5 µl	5 µl	5 µl	
1x PARP buffer	20 µl	-	-	-
TNKS2 (~ 0.75 ng/µl)	–	20 µl	20 µl	20 µl
Total	50 µl	50 µl	50 µl	50 µl


- 6) Add 5 µl of inhibitor solution to each well designated “Test Inhibitor”. For the “Positive Control”, “Substrate Control”, and “Blank”, add 5 µl of the same solution without inhibitor (inhibitor buffer).
- 7) Add 20 µl of 1x PARP buffer to the well designated “Blank”.
- 8) Initiate the reactions by adding 20 µl of diluted TNKS2 prepared as described above. Incubate the reactions for 1 hour at room temperature.
- 9) Wash the plate 3 times with 200 µl PBST per well.

Step 3: Detection

- 1) Dilute **Streptavidin-HRP** 1:50 in Blocking buffer.
- 2) Add 50 µl of diluted **Streptavidin-HRP** to each well. Incubate for 30 minutes at room temperature.
- 3) Wash three times with 200 µl PBST buffer as above.
- 4) Just before use, mix on ice 50 µl **HRP chemiluminescent substrate A** and 50 µl **HRP chemiluminescent substrate B** and add 100 µl per well. Discard any unused chemiluminescent reagent after use.
- 5) Immediately read sample in a luminometer or microtiter-plate capable of reading chemiluminescence.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

Example of Assay Results:




Inhibition of TNKS2 enzyme (#80515) with XAV939 #27100), measured using the *TNKS2 Histone Ribosylation Assay Kit (Biotin-labeled NAD⁺)*, (Catalog # 80578). Luminescence was measured using a Bio-Tek fluorescent microplate reader.

RELATED PRODUCTS:

<u>Product</u>	<u>Cat. #</u>	<u>Size</u>
PARP2 Assay Kit	#80552	96 rxns.
PARP3 Assay Kit	#80553	96 rxns.
TNKS1 Histone Ribosylation Assay Kit (Antibody Detection)	#80574	96 rxns.
PARP5b (TNKS2) Assay Kit	#80576	96 rxns.
PARP6 Assay Kit	#80556	32 rxns.
PARP1 Enzyme	#80501	10 µg
PARP2 Enzyme	#80502	10 µg
PARP3 Enzyme	#80503	10 µg
PARP6 Enzyme	#80506	10 µg
TNKS1 (PARP5A) Enzyme	#80504	10 µg
TNKS2 (PARP5B), (667-end) Enzyme	#80505	10 µg
TNKS2 (PARP5B), (849-end) Enzyme	#80515	10 µg
PARP7 Enzyme	#80507	10 µg
PARP9 Enzyme	#80509	10 µg
PARP11 Enzyme	#80511	10 µg
PARP12 Enzyme	#80512	10 µg

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



UK & Rest of World
184 Milton Park, Abingdon
OX14 4SE, Oxon, UK
Tel: +44 (0) 1235 828 200
Fax: +44 (0) 1235 820 482

Switzerland
Centro Nord-Sud 2E
CH-6934 Bioggio-Lugano
Tel: +41 (0) 91 604 55 22
Fax: +41 (0) 91 605 17 85

Deutschland
Bockenheimer Landstr. 17/19
60325 Frankfurt/Main
Tel: +49 (0) 69 779099
Fax: +49 (0) 69 13376880

North America
23591 El Toro Rd, Suite #180
Lake Forest, CA 92630
Tel: +1 800 987 0985
Fax: +1 949 265 7703

amsbio
info@amsbio.com
www.amsbio.com
AMS Biotechnology