


Synonym

ACE-2,ACEH,ACE2

Source

Human ACE2, His Tag (AMS.AC2-H52H8) is expressed from human 293 cells (HEK293). It contains AA Gln 18 - Ser 740 (Accession # Q9BYF1-1). Predicted N-terminus: His

Molecular Characterization

This protein carries a polyhistidine tag at the N-terminus.

The protein has a calculated MW of 85.5 kDa. The protein migrates as 100-115 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Delivered as bulk protein in a 0.2 µm filtered solution of 50 mM Tris, 150 mM NaCl, pH7.5 with glycerol as protectant.

Contact us for customized product form or formulation.

Storage


Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- The product MUST be stored at -70°C or lower upon receipt;
- -70°C for 3 months under sterile conditions.

Shipping

This product is supplied as sterile liquid solution and shipped frozen with dry ice, please inquire the shipping cost.

SDS-PAGE



Human ACE2, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA**SEC-MALS**

The purity of Human ACE2, His Tag (Cat. No. AMS.AC2-H52H8) was more than 90% and around 190-220 kDa verified by SEC-MALS.

Human ACE2 / ACEH Protein, His Tag (MALS verified)


Catalog # AMS.AC2-H52H8



Immobilized SARS-CoV-2 S protein RBD, Fc Tag at 2µg/mL (100 µL/well) can bind Human ACE2, His Tag with a linear range of 0.04-0.313 µg/mL (QC tested).

Immobilized Human ACE2, His Tag at 2 µg/mL (100µL/well) can bind Biotinylated SARS-CoV-2 S1 protein, His,Avitag with a linear range of 0.02-0.156 µg/mL (Routinely tested).

Bioactivity-SPR



SARS-CoV-2 S1 protein, Fc Tag (Cat. No. AMS.S1N-C5255) captured on CM5 Chip via anti-human IgG Fc antibodies can bind Human ACE2, His Tag (Cat. No. AMS.AC2-H52H8) with an affinity constant of 3.79 nM as determined in SPR assay (Biacore T200) (Routinely tested).

Background

Angiotensin-converting enzyme 2 (ACE2) is also known as ACEH (ACE homolog), is an integral membrane protein with considerable homology to ACE, which belongs to the peptidase M2 family. ACE2 is an exopeptidase that catalyses the conversion of angiotensin I to the nonapeptide angiotensin, or the conversion of angiotensin II to angiotensin 1-7. ACE2 may be an important regulator of heart function. In case of human coronaviruses SARS and HCoV-NL63 infections, ACE-2 serve as functional receptor for the spike glycoprotein of both coronaviruses. ACE2 is activated by chloride and fluoride, but not bromide and Inhibited by MLN-4760, cFP_Leu, and EDTA, but not by the ACE inhibitors lisinopril, captopril and enalaprilat. ACE2 is active from pH 6 to 9, and the optimum pH is 6.5 in the presence of 1 M NaCl.

References

- (1) [Turner AJ, et al., 2002, Can. J. Physiol. Pharmacol. 80\(4\): 346-53.](#)
- (2) [Katovich MJ, et al., 2005, Exp. Physiol. 90\(3\): 299-305.](#)
- (3) [Donoghue M, et al., 2000, Circ. Res. 87:E1-E9.](#)
- (4) [Tipnis S.R, et al., 2000, J. Biol. Chem. 275:33238-33243.](#)

AMSBIO | www.amsbio.com | info@amsbio.com

 **UK & Rest of the World**
184 Park Drive, Milton Park
Abingdon OX14 4SE, UK
T: +44 (0)1235 828 200
F: +44 (0) 1235 820 482

 **North America**
1035 Cambridge Street,
Cambridge, MA 02141
T: +1 (617) 945-5033 or
T: +1 (800) 987-0985
F: +1 (617) 945-8218

 **Germany**
Bockenheimer Landstr. 17/19
60325 Frankfurt/Main
T: +49 (0) 69 779099
F: +49 (0) 69 13376880

 **Switzerland**
Centro Nord-Sud 2E
CH-6934 Bioggio-Lugano
T: +41(0) 91 604 55 22
F: +41(0) 91 605 17 85