

**TurboCells™ BL21(DE3)
TurboCells™ BL21(DE3)pLysS
Chemically Competent *E. coli***

Instruction Manual

Catalog Numbers

**AMS.C302020
AMS.C303020**

AMSBIO | www.amsbio.com | info@amsbio.com



UK & Rest of the World
184 Park Drive, Milton Park
Abingdon OX14 4SE, UK
T: +44 (0)1235 828 200
F: +44 (0) 1235 820 482



North America
1035 Cambridge Street,
Cambridge, MA 02141
T: +1 (617) 945-5033 or
T: +1 (800) 987-0985
F: +1 (617) 945-8218



Germany
Bockenheimer Landstr. 17/19
60325 Frankfurt/Main
T: +49 (0) 69 779099
F: +49 (0) 69 13376880



Switzerland
Centro Nord-Sud 2E
CH-6934 Bioggio-Lugano
T: +41(0) 91 604 55 22
F: +41(0) 91 605 17 85

Purchaser Notification

Limited License

The purchase price paid for the TurboCells™ BL21(DE3) Chemically Competent *E. coli* and TurboCells™ BL21(DE3)pLysS Chemically Competent *E. coli* by end users grants them a non-transferable, non-exclusive license to use the kit and/or its separate and included components (as listed in the Kit Contents section). This kit is intended **for internal research use only** by the purchaser. Furthermore, **research use only** means that this kit and all of its contents are excluded, without limitation, from resale, repackaging, or use for the making or selling of any commercial product or service without the written approval of Genlantis.

Academic and Non-Profit Laboratory Use

The T7 expression system is based on technology developed at Brookhaven National Laboratory under contract with the U. S. Department of Energy and is the subject of patent applications assigned to Brookhaven Science Associates, LLC. (BSA). BSA will grant a non-exclusive license for use of this technology, including the enclosed materials, based upon the following assurances:

1. These materials are to be used for noncommercial research purposes only. A separate license is required for any commercial use, including the use of these materials for research purposes or production purposes by any commercial entity. Information about commercial licenses may be obtained from: The Office of Intellectual Property & Sponsored Research, Brookhaven National Laboratory, Bldg. 475D, P. O. Box 5000, Upton, New York 11973-5000, telephone (631) 344-7134.
2. No materials that contain the cloned copy of T7 gene l, the gene for T7 RNA polymerase, may be distributed further to third parties outside of your laboratory, unless the recipient receives a copy of this license and agrees to be bound by its terms. This limitation applies to strains BL21(DE3), BL21(DE3)pLysS, and BL21(DE3)pLysE, and any derivatives you may make of them.

You may refuse this license by returning the enclosed materials unused. By keeping or using the enclosed materials, you agree to be bound by the terms of this license.

Limited Warranty

Genlantis is committed to providing its customers with high-quality products and services. Our goal is to ensure that every customer is 100% satisfied. If you should have any questions or concerns about this product, please contact our technical service department. Genlantis warrants that all of its products will perform according to the specifications stated on the certificate of analysis. The company will replace, free of charge, any product that does not meet those specifications. This warranty limits Genlantis' liability only to the cost of the product. No warranty is granted for products beyond their listed expiration date. No warranty is applicable unless all product components are stored in accordance with instructions. Genlantis reserves the right to select the method(s) used to analyze a product unless Genlantis agrees to a specified method in writing prior to acceptance of the order.

Product Use Limitations

The TurboCells™ BL21(DE3) Chemically Competent *E. coli* and TurboCells™ BL21(DE3)pLysS Chemically Competent *E. coli* and all other components are developed, designed, intended, and sold for research use only. They are not to be used for human diagnostic or included/used in any drug intended for human use. All care and attention should be exercised in the handling of the kit components by following appropriate research lab practices.

Kit Contents and Ordering Information

The TurboCells™ BL21(DE3) Chemically Competent *E. coli* and TurboCells™ BL21(DE3)pLysS Chemically Competent *E. coli* kit contains sufficient competent cells for 20 transformations.

Catalog Number	Contents	Amount
C302020	TurboCells™ BL21(DE3) Chemically Competent <i>E. coli</i>	20 x 50 µl
	TurboCells™ BL21(DE3) Genotype: F ⁻ <i>ompT hsdS_B (r_B⁻ m_B⁻) gal dcm (DE3)</i>	
	TurboCells™ Transformation Buffer	5 ml
	pUC19 Positive Control Plasmid <i>Provided as a positive control to verify the transformation efficiency of the competent cells.</i>	20 µl (500 pg/µl)
C303020	TurboCells™ (DE3)pLysS Chemically Competent <i>E. coli</i>	20 x 50 µl
	TurboCells™ (DE3)pLysS Genotype: F ⁻ <i>ompT hsdSB (r_B⁻ m_B⁻) gal dcm (DE3) pLysS (Cam^R)</i>	
	TurboCells™ Transformation Buffer	5 ml
	pUC19 Positive Control Plasmid. <i>Provided as a positive control to verify the transformation efficiency of the competent cells.</i>	20 µl (500 pg/µl)

Shipping and Storage

The TurboCells™ BL21(DE3) Chemically Competent *E. coli* and TurboCells™ BL21(DE3)pLysS Chemically Competent *E. coli* are shipped frozen. For maximum stability and long-term use, store cells at -70°C upon receipt. The TurboCells™ Transformation Buffer should be stored at room temperature. All components are stable for six months when stored properly.

Related Products

For convenient, safe, and optimal plating of bacterial cells on standard agar plates:

Product Name	Cat. No.	Quantity
EZ-Spread™ Plating Beads, Single-Use Tubes	C400050	50 tubes
EZ-Spread™ Plating Beads, Dispenser Bottle	C400100	1 bottle/100 plates

Introduction

TurboCells™ BL21(DE3) and TurboCells™ BL21(DE3)pLysS Chemically Competent *E. coli* incorporate a unique 3-minute transformation protocol with efficient transgene expression from T7 promoter-based vectors. Both strains contain the λ DE3 lysogen, which expresses T7 RNA polymerase under the control of a *lacUV5* promoter. Upon addition of IPTG, the expression of the T7 RNA polymerase is induced, which allows for high-level protein expression from T7 promoter-based expression vectors (e.g., pET). Additionally, BL21(DE3)pLysS contains the pLysS plasmid which produces T7 lysozyme, an inhibitor of T7 RNA polymerase, to reduce basal level expression of proteins that are toxic in *E. coli*.

Unlike other commercially available BL21 strains, the TurboCells™ BL21(DE3) and TurboCells™ BL21(DE3)pLysS allow efficient transformation without the need for a lengthy transformation reaction or hour-long recovery after heat shock. In addition, with TurboCells™, you can perform the heat shock step at 37°C, eliminating the inconvenient 42°C water bath. TurboCells™ Competent *E. coli* are great for routine protein expression experiments. The fast and easy protocol also makes them ideal for high-throughput applications.

When using the 3-minute transformation protocol, you can achieve transformation efficiencies of up to 5×10^5 cfu/ μ g of supercoiled DNA, which is more than sufficient for most protein expression experiments. When using the traditional 1.5 – 2 hour transformation protocol with the TurboCells™ BL21(DE3) and TurboCells™ BL21(DE3)pLysS Competent *E. coli*, greater transformation efficiencies can be achieved.

For optimal transformation results and your convenience, TurboCells™ BL21(DE3) and TurboCells™ BL21(DE3)pLysS Chemically Competent *E. coli* are supplied in single-use 50 μ l aliquots. This format avoids efficiency-robbing freeze-thaw cycles and wasted cells.

Table 1. Genotypes of TurboCells™ Chemically Competent *E. coli*

TurboCells™ BL21(D3): F ⁻ <i>ompT hsdS_B</i> (<i>r_B⁻ m_B⁻</i>) <i>gal dcm</i> (DE3)	
TurboCells™ BL21(D3)pLysS: F ⁻ <i>ompT hsdS_B</i> (<i>r_B⁻ m_B⁻</i>) <i>gal dcm</i> (DE3) pLysS (Cam ^R)	
Genotype	Advantage
<i>DE3</i>	Encodes T7 lysogen for T7 RNA polymerase for high-level transcription.
<i>pLysS</i>	Reduced basal expression of the T7 RNA polymerase. This reduces basal expression of toxic transgenes.
<i>Cam^R</i>	Chloramphenicol resistance; enables maintenance of the pLysS plasmid
<i>ompT</i>	Deficient in the OmpT protease, resulting in a higher yield of intact recombinant proteins.
<i>hsd SB</i> (<i>r_B⁻ m_B⁻</i>)	Improved cloning efficiencies and representations of methylated DNA

1. Transformation

- 1.1 Thaw one tube of the TurboCells™ Chemically Competent *E. coli* on ice.
- 1.2 Transfer to pre-chilled Falcon 2059 tube.
- 1.3 Add 5 to 50 ng of ligation reaction, in a volume of 1 to 10 µl (or add 1 µl of 500 pg/µl of pUC19 control vector) to the cells; mix by tapping gently and incubate on ice for 3 to 5 minutes.
- 1.4 Heat shock the mix at 37°C for 60 seconds.
- 1.5 Place on ice immediately and incubate for 2 minutes. (Recommended)
- 1.6 Dilute the transformation reaction with 200 µl TurboCells™ Transformation Buffer and spread 25–100 µl of transformed cells on 37°C pre-warmed LB/Agar plates containing the appropriate antibiotic selection (e.g. 50 µg/ml ampicillin). We recommend using EZ-Spread™ Plating Beads (Cat. No. C400050 & C400100) to obtain optimal plating results. If using <100 µl of transformed cells, add enough TurboCells™ Transformation Buffer to make up 100 µl for ease of spreading.

IMPORTANT

If kanamycin is used for antibiotic selection, add 0.25 ml room temperature SOC medium and incubate at 37°C with horizontal shaking at 225 rpm for 1 hour in an air incubator, prior to adding 600 µl TurboCells™ Transformation Buffer.

- 1.7 Incubate overnight at 37°C.

Notes:

- I. *If maximum numbers of colonies are desired, replace Step 4 with the following: Collect cells by spinning in a microfuge for 10 seconds. Resuspend cell pellet in 50 µl TurboCells™ Transformation Buffer and spread on an agar plate containing antibiotic.*
- II. *If satellite colonies appear on the plates after overnight incubation, we recommend the following: 1) reduce the volume of transformation reaction used for plating, 2) if ampicillin is used, replace it with a more stable homologue such as Carbenicillin (disodium salt), or 3) increase the antibiotic concentration in the plate.*
- III. *It is not necessary to dilute your ligation mix with TE. The TurboCells™ BL21(DE3) and BL21(DE3)pLysS competent cells are specially prepared to work with most ligation buffers. In our tests, up to 10 µl of undiluted ligation mix could be used without significantly compromising the transformation efficiency.*
- IV. *To increase transformation efficiencies and obtain higher number of colonies, we recommend the following traditional protocol:*
 1. *Thaw one tube of the TurboCells™ competent cells on ice (10-15 minutes).*
 2. *Add 1-10 µl of ligation mix to the cells; mix gently and incubate on ice for 15 to 30 minutes.*
 3. *Heat the mix at 42°C for 45 seconds.*

4. Add 0.25 ml room temperature SOC medium and incubate at 37°C for 1 hour in an air incubator. Shaking tubes horizontally at 225 rpm is recommended for the best efficiency.
 5. Dilute the transformation reaction if necessary and spread 100 µl of transformed cells on LB/Agar plates containing the appropriate antibiotic selection (e.g. ampicillin or kanamycin).
 6. Incubate overnight at 37°C.
- V. Transformation efficiencies for ligation of inserts to vectors will be slightly lower than supercoiled plasmid (between 5 to 10 fold lower).
- VI. To test the efficiency of competent cells using the provided supercoiled pUC19 plasmid DNA, please use the following protocol:
1. Transform 1 µl (500 pg) pUC19 into 50 µl of competent cells. Incubate on ice for 3 minutes.
 2. Heat shock the mix at 37°C for 60 seconds. Place on ice immediately and incubate for 2 minutes.
 3. Dilute the transformation reaction 10 fold with TurboCells™ Transformation Buffer and plate 50 µl on a LB agar plate containing 50 µg/ml ampicillin.
 4. Incubate overnight at 37°C and count colonies. Calculate transformation efficiency as follows:

$$\frac{\text{Number of colonies}}{500 \text{ pg pUC19}} \times \frac{1 \times 10^6 \text{ pg}}{\mu\text{g}} \times \frac{50 \mu\text{l}}{50 \mu\text{l plated}} \times 10 \text{ (dilution factor)} = \text{CFU} / \mu\text{g}$$

The expected control efficiency = 5×10^5 CFU/µg.

2. Example Protocol for Protein Expression

- 2.1 Following transformation, pick 3 or 4 individual colonies, and with each one, inoculate 1 - 50 ml aliquots of LB medium containing the appropriate antibiotic to select for the expression vector.

IMPORTANT If using TurboCells™ BL21(DE3)pLysS, include 40 µg/ml chloramphenicol in the LB medium to maintain the pLysS plasmid.

TIP Perform a fresh transformation of BL21(DE3)pLysS cells before each induction experiment.

- 2.2 Incubate each aliquot overnight with shaking at 37°C.
- 2.3 The following day, transfer a 50 µl aliquot of each culture to 1 ml of fresh LB medium without antibiotics, and incubate for approximately 2 hours at 37°C until OD₆₀₀ reaches 0.4–0.9.
- 2.4 Remove a 100 µl aliquot of each culture and place on ice as a non-induced negative control.

- 2.5 With the remaining 950 µl of each culture, induce by adding IPTG to a final concentration of 1.0 mM, and culture for 2-3 hours. A range of IPTG concentrations (0.4 mM – 1.0 mM) and time points is recommended to determine optimal conditions for expression of your protein.
- 2.6 After the end of the induction period, protein expression may be analyzed by Western blot, SDS-PAGE, enzymatic assay, *etc.*
- 2.7 It is possible to save the transformed TurboCells™ BL21(DE3) cells if desired. Simply isolate the cells by centrifugation at 5000 x g for 5 minutes. Remove the supernatant and store the cells as a frozen pellet at -70°C.

Notes:

- I. *The above section provides general guidelines for performing protein expression in TurboCells™ BL21(DE3) and TurboCells™ BL21(DE3)pLysS Chemically Competent E. coli using T7 RNA polymerase-based expression plasmids. These guidelines are **optional**, and they may be modified or discarded as required by your particular application.*
- II. *For expression of proteins known to be **non-toxic** in E. coli, we recommend using the TurboCells™ BL21(DE3), as this strain provides the highest levels of protein expression in most cases.*
- III. *For expression of proteins known to be **toxic** in E. coli, we recommend using the TurboCells™ BL21(DE3)pLysS strain. This strain contains the pLysS plasmid, which encodes T7 lysozyme. T7 RNA polymerase is significantly inhibited by T7 lysozyme, thus “leakage” (i.e., basal level expression) of toxic recombinant proteins in uninduced cells is significantly reduced.*
- IV. *We recommend that TurboCells™ BL21(DE3) and TurboCells™ BL21(DE3)pLysS be used **for expression only**. To maintain and propagate your T7 expression vector, use an E. coli strain that does not contain the T7 RNA polymerase, e.g. TurboCells™ Chemically Competent E. coli.*

Product Name	Cat. No.	Quantity
TurboCells™ Chemically Competent <i>E. coli</i>	C300020	20 x 50 µl
TurboCells™ F' Chemically Competent <i>E. coli</i>	C301020	20 x 50 µl

Quality Control

Kit Component	Quality Control Standard
TurboCells™ BL21(DE3) and TurboCells™ BL21(DE3)pLysS Chemically Competent <i>E. coli</i>	Yield 5×10^5 cfu/ μ g transformation efficiency when transformed with pUC19 plasmid using the above protocol.

Troubleshooting Guide

Problem	Possible Causes	Recommended Solutions
No or very few colonies on agar plate after transformation	Gene of interest is toxic in <i>E. coli</i>	Use TurboCells™ BL21(DE3)pLysS Competent <i>E. coli</i> to reduce basal level protein expression.
	Competent cells are dead due to improper storage or shipping conditions	The efficiency of competent cells can be verified using a plasmid such as pUC19. Make sure you store the SmartCells™ Competent <i>E. coli</i> at -70°C .
	Wrong antibiotic selection.	Double-check that correct antibiotic selection was used.
No or very little protein expression detected after induction.	Poor vector design	Check if expression cassette is in-frame with promoter.
	Induction time and IPTG concentration not optimized.	Optimize IPTG to a final concentration of 0.4 – 1.0 mM, and culture time for 2-3 hours. Also, temperature may be optimized between $25-37^{\circ}\text{C}$.

AMSBIO | www.amsbio.com | info@amsbio.com

 **UK & Rest of the World**
184 Park Drive, Milton Park
Abingdon OX14 4SE, UK
T: +44 (0)1235 828 200
F: +44 (0) 1235 820 482

 **North America**
1035 Cambridge Street,
Cambridge, MA 02141
T: +1 (617) 945-5033 or
T: +1 (800) 987-0985
F: +1 (617) 945-8218

 **Germany**
Bockenheimer Landstr. 17/19
60325 Frankfurt/Main
T: +49 (0) 69 779099
F: +49 (0) 69 13376880

 **Switzerland**
Centro Nord-Sud 2E
CH-6934 Bioggio-Lugano
T: +41(0) 91 604 55 22
F: +41(0) 91 605 17 85