amsbio» Accelerate Your Discovery

EZ-Spread[™] Plating Beads

Cat. No. AMS.C400050 Single-Use Tubes Cat. No. AMS.C400100 Dispenser Bottle

Product Summary

Cat. No: AMS.C400050

Description: EZ-Spread[™] Plating Beads Single-Use Tubes (Sterile). Each box contains 50 tubes for 50 platings.

Cat. No: AMS.C400100

Description: EZ-Spread[™] Plating Beads Dispenser Bottle (Sterile). Each bottle contains sufficient beads for approximately 100 standard 100 mm agar plates.

Store both products at room temperature.

INTRODUCTION

EZ-SpreadTM Plating Beads are specifically designed for convenient, safe, and optimal plating of bacterial and yeast cells on standard agar plates. The EZ-SpreadTM glass beads have been manufactured using special molds to give consistent shape and diameter that yield optimal plating efficiencies, often times exceeding the traditional method using bent glass rods The EZ-Spread[™] Plating Beads offer the following benefits:

- Sterile plating in just seconds
- No need for glass rods and potentially hazardous ethanol flaming.
- More uniform spreading and more colonies
- Two choices of ready-to-use sterile packaging-dispenser bottle for spreading 100 plates or single-use tubes for 50 plating
- Fastest and easiest way to spread cells on multiple plates simultaneously

PROTOCOLS

EZ-Spread[™] Glass Beads greatly simplify the process of plating cells making them faster, safer, easier, and more convenient. They are provided sterile and ready-to-use. For best results, use the following instructions.

For Single-Use Tubes

- 1. Gather materials under laminar flow hood. (Optional)
- 2. Remove needed number of tubes from box and lightly spray exterior of tubes with 70% ethanol solution.
- 3. Pipette 50 to 400 µl of transformed cells onto 100 mm Agar plate.
- 4. Open one tube and transfer EZ-Spread[™] Plating Beads onto agar surface. Each tube contains approximately 20 sterile EZ-Spread[™] Glass Beads. Discard empty tubes when done.

AMSBIO | www.amsbio.com | info@amsbio.com

AMSBIO LLC USA & Canada	AMSBIO Europe BV EU
1035 Cambridge Street,	Berenkoog 41,
Cambridge, MA 02141	1822 BH Älkmaar,
T: +1 (617) 945-5033 or	Netherlands
T: +1 (800) 987-0985	T: +31 (0) 72 808024

F: +1 (617) 945-8218

erenkoog 41, 22 BH Alkmaar, etherlands +31 (0) 72 8080244 F: +31 (0) 72 8080142

UK & Rest of the World 184 Park Drive, Milton Park

AMS Biotechnology (Europe) Ltd

Abingdon OX14 4SE T: +44 (0) 1235 828 200 F: +44 (0) 1235 820 482 AMS Biotechnology (Europe) Ltd Switzerland

Via Lisano 3,

CH-6900 Massagno

T: +41 (0) 91 604 55 22

F: +41 (0) 91 605 17 85

(CP.683)

AMSBIO Europe BV Deutschland

T: +49 (0) 69 779099 F: +49 (0) 69 13376880

- 5. Cover the plate and roll the beads around in a circular motion for about 30 seconds, or until the surface of the agar plate seems dry. If using more than one plate, stack covered plates up and roll the whole stack around with vigorous motion. The aim is to move the beads around the plates as evenly as possible so that the transformed cells spread uniformly across the entire surface of the agar plate.
- 6. Remove the lid(s) from the plate(s) and flip upside down to discard the beads into an appropriate hazardous waste container.

For dispenser bottles

- 1. Gather materials under laminar flow hood. (Optional)
- 2. Remove shrink-wrap from the lid and lightly spray with 70% ethanol solution; wipe with a napkin or tissue until dry.
- 3. Pipette 50 to 400 µl of transformed cells onto 100 mm agar plate.
- 4. Flip up desired side of lid for either bulk or slow dispensing of beads.
- 5. Tilt bottle upside down until approximately 20 beads are dispensed onto a standard 100 mm plate. Close dispenser lids to maintain sterility of contents as soon as finished with dispensing to maintain sterility of contents.
- 6. Cover the plate and roll the beads in a circular motion for about 30 seconds, or until the surface of the agar plate seems dry. If using more than one plate, stack covered plates up and roll the whole stack around with vigorous motion. The aim is to move the beads around the plates as evenly as possible so that the transformed cells spread uniformly across the entire surface of the agar plate.
- 7. Remove the lid(s) from the plate(s) and flip upside down to discard the beads into an appropriate hazardous waste container.

AMSBIO | www.amsbio.com | info@amsbio.com

AMSBIO LLC USA & Canada

1035 Cambridge Street, Cambridge, MA 02141 T: +1 (617) 945-5033 or T: +1 (800) 987-0985 F: +1 (617) 945-8218



Berenkoog 41, 1822 BH Alkmaar, Netherlands T: +31 (0) 72 8080244 F: +31 (0) 72 8080142 AMS Biotechnology (Europe) Ltd UK & Rest of the World

184 Park Drive, Milton Park Abingdon OX14 4SE T: +44 (0) 1235 828 200 F: +44 (0) 1235 820 482 AMS Biotechnology (Europe) Ltd Switzerland

Via Lisano 3, (CP.683) CH-6900 Massagno T: +41 (0) 91 604 55 22 F: +41 (0) 91 605 17 85 AMSBIO Europe BV Deutschland

T: +49 (0) 69 779099 F: +49 (0) 69 13376880