

Hacat Cell Line

Catalog No.: AMS.CL-0090

Origin and General Characteristics

Cell Name	Hacat Cell Line
Age	62 years old
Tissue	Skin
Morphology	Epithelial
Growth Properties	Adherent
Descriptions	In vitro spontaneously transformed keratinocytes from histologically normal skin.
Biosafety Level	1

Culture Conditions and Handling

Complete Growth Medium	MEM +15% FBS +1% P/S
Subculturing	Remove and discard culture medium. Briefly rinse the cell layer with DPBS solution to remove all traces of serum that contains trypsin inhibitor. Add 1.0 to 2.0 mL of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 2 to 3 minutes). Cells that are difficult to detach may be placed at 37°C to facilitate dispersal. Add 4.0 to 6.0 mL of complete growth medium and aspirate cells by gently pipetting. Add appropriate aliquots of the cell suspension to new culture vessels.
Subcultivation Ratio	1:2-1:4
Medium Renewal	Every 2 to 3 days
Cryopreservation	Freeze medium: 60% Basal medium+30% FBS+10% DMSO Storage temperature: Liquid nitrogen vapor phase
Culture Conditions	Atmosphere: Air, 95%; CO ₂ , 5% Temperature: 37°C

Special Features of the Cell Line

Tumorigenic	No
--------------------	----

Recommendations for handling of cryopreserved cells

- The cell is packaged by dry ice. When receiving the cell, please make sure that the vial is still frozen. If there is cell thawing in the tube, please take photo before experiment or storage.
- If immediate culturing is not intended, the cryovial(s) must be stored in liquid nitrogen (-196°C) or at least at -80°C after arrival.
If immediate culturing is intended, please follow these instructions:
- Quickly thaw by rapid agitation in a 37°C water bath within 45-90 seconds. The water bath should have clean water containing an antimicrobial agent. As soon as the sample has thawed, remove the cryovial from the water bath.
Note: A small ice clump should still remain and the vial should still be cold.
From now on, all operations should be carried out under aseptic conditions.
- Transfer the cryovial to a sterile flow cabinet and wipe with 70% alcohol. Carefully open the vial and transfer the cell suspension into a 15 ml centrifuge tube containing 9 ml of cell complete medium (room temperature or 37°C).

AMSBIO | www.amsbio.com | info@amsbio.com



UK & Rest of the World
184 Park Drive, Milton Park
Abingdon OX14 4SE, UK
T: +44 (0)1235 828 200
F: +44 (0) 1235 820 482



North America
1035 Cambridge Street,
Cambridge, MA 02141
T: +1 (617) 945-5033 or
T: +1 (800) 987-0985
F: +1 (617) 945-8218



Germany
Bockenheimer Landstr. 17/19
60325 Frankfurt/Main
T: +49 (0) 69 779099
F: +49 (0) 69 13376880



Switzerland
Centro Nord-Sud 2E
CH-6934 Bioggio-Lugano
T: +41(0) 91 604 55 22
F: +41(0) 91 605 17 85

5. In order to reduce cell damage, add 1ml of cell complete medium into cryovial, slightly pipette, then use a pipettor to add 1 ml of suspension into the centrifuge tube. Resuspend the cells carefully. Centrifuge at 300×g for 3 min and discard the supernatant. The centrifugation step may be omitted, but in this case the remains of the freeze medium have to be removed 24 hours later.
6. Resuspend the cells carefully in 10ml fresh cell culture medium and transfer them into one or two T25 cell culture flasks.

AMSBIO | www.amsbio.com | info@amsbio.com



UK & Rest of the World
184 Park Drive, Milton Park
Abingdon OX14 4SE, UK
T: +44 (0)1235 828 200
F: +44 (0) 1235 820 482



North America
1035 Cambridge Street,
Cambridge, MA 02141
T: +1 (617) 945-5033 or
T: +1 (800) 987-0985
F: +1 (617) 945-8218



Germany
Bockenheimer Landstr. 17/19
60325 Frankfurt/Main
T: +49 (0) 69 779099
F: +49 (0) 69 13376880



Switzerland
Centro Nord-Sud 2E
CH-6934 Bioggio-Lugano
T: +41(0) 91 604 55 22
F: +41(0) 91 605 17 85