

Synonym

S protein RBD, Spike glycoprotein Receptor-binding domain, S glycoprotein RBD, Spike protein RBD, COVID-19

Source

SARS-CoV-2 S protein RBD, His Tag (AMS.SPD-C52H3) is expressed from human293 cells (HEK293). It contains AA Arg 319 - Lys 537 (Accession # [QHD43416.1](#)). Predicted N-terminus: Arg 319

Molecular Characterization

S protein RBD(Arg 319 - Lys 537) QHD43416.1	Poly-his
--	----------

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 26.5 kDa. The protein migrates as 32-35 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4. Normally trehalose is added as protectant before lyophilization.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE


SARS-CoV-2 S protein RBD, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA**SEC-MALS**


The purity of SARS-CoV-2 S protein RBD, His Tag (Cat. No. AMS.SPD-C52H3) was more than 90% and the molecular weight of this protein is around 30-40 kDa verified by SEC-MALS.

SARS-CoV-2 (COVID-19) S protein RBD, His Tag (MALS verified)


Catalog # AMS.SPD-C52H3



Immobilized Human ACE2, Fc Tag at 2 µg/mL (100µL/well) can bind SARS-CoV-2 S protein RBD, His Tag with a linear range of 8-125 ng/mL (QC tested).




Immobilized SARS-CoV-2 S protein RBD, His Tag at 1 µg/mL (100 µL/well) can bind Human ACE2, Fc Tag with a linear range of 0.2-3 ng/mL (Routinely tested).



Immobilized SARS-CoV-2 S protein RBD, His Tag at 1 µg/mL (100 µL/well) can bind Anti-SARS-CoV-2 Neutralizing Antibody, Human IgG1 with a linear range of 0.1-3 ng/mL (Routinely tested).

Bioactivity-SPR




Human ACE2, Fc Tag (Cat. No. AMS.AC2-H5257) captured on CM5 chip via anti-human IgG Fc antibodies surface can bind SARS-CoV-2 S protein RBD, HisTag (Cat. No. AMS.SPD-C52H3) with an affinity constant of 17 nM as determined in a SPR assay (Biacore T200) (Routinely tested).

SARS-CoV-2 (COVID-19) S protein RBD, His Tag (MALS verified)

Catalog # AMS.SPD-C52H3

Bioactivity-BLI



Loaded Human ACE2, Fc Tag (Cat. No. AMS.AC2-H5257) on Protein A Biosensor, can bind SARS-CoV-2 S protein RBD, His Tag (Cat. No. AMS.SPD-C52H3) with an affinity constant of 34.5 nM as determined in BLI assay (ForteBio OctetRed96e) (Routinely tested).

Background

It's been reported that SARS-CoV-2 can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

References

- (1) [Wan Y, et al. J Virol. 2020. pii: JVI.00127-20.](#)
- (2) [Benvenuto D, et al. J Med Virol. 2020.](#)
- (3) [Chang CY, et al. AMB Express. 2020. 10\(1\):20.](#)

AMSBIO | www.amsbio.com | info@amsbio.com

 **UK & Rest of the World**
184 Park Drive, Milton Park
Abingdon OX14 4SE, UK
T: +44 (0) 1235 828 200
F: +44 (0) 1235 820 482

 **North America**
1035 Cambridge Street,
Cambridge, MA 02141
T: +1 (617) 945-5033 or
T: +1 (800) 987-0985
F: +1 (617) 945-8218

 **Germany**
Bockenheimer Landstr. 17/19
60325 Frankfurt/Main
T: +49 (0) 69 779099
F: +49 (0) 69 13376880

 **Switzerland**
Centro Nord-Sud 2E
CH-6934 Bioggio-Lugano
T: +41(0) 91 604 55 22
F: +41(0) 91 605 17 85