


Mouse Anti-Human CD44 Std. Monoclonal Antibody (Clone SPM521)

CATALOG #:	E17360 0.1 ml mouse monoclonal antibody purified by protein A/G in PBS/1% BSA buffer pH 7.6 with less than 0.1% sodium azide.
	E17364 1 ml mouse monoclonal antibody purified by protein A/G in PBS/1% BSA buffer pH 7.6 with less than 0.1% sodium azide.
	E17361 7 ml pre-diluted mouse monoclonal antibody purified by protein A/G in TBS/1% BSA buffer pH 7.6 with less than 0.1% sodium azide. <i>(For IHC only)</i>



Human esophageal carcinoma stained with anti-CD44 antibody

INTENDED USE:	For Research Use Only. Not for use in diagnostic procedures.
CLONE:	SPM 521
IMMUNOGEN:	A synthetic peptide corresponding to human CD44.
IG ISOTYPE:	IgG _{2a}
EPITOPE:	Internal region
MOLECULAR WEIGHT	80kDa
SPECIES REACTIVITY:	Human (tested). Others not tested.
DESCRIPTION:	CD44 is a cell surface glycoprotein expressed on lymphocytes, monocytes, and granulocytes. CD44 is also known as homing cell adhesion molecule (H-CAM), phagocytic glycoprotein-1 (PgP-1), ECM-III, HUTCH-1, or Hermes-1. CD44 has been implicated in cell migration, in lymphocyte homing, and in tumor metastasis.
APPLICATIONS:	Immunohistochemistry (IHC) and Western Blotting
IHC PROCEDURE:	<p>Specimen Preparation: Formalin-fixed, paraffin-embedded tissues are suitable for use with this primary antibody.</p> <p>Deparaffinization: Deparaffinize slides using xylene or xylene alternative and graded alcohols.</p> <p>Antibody Dilution: If using the concentrate format of this product, dilute the antibody 1:100. The dilutions are estimates; actual results may differ because of variability in methods and protocols.</p> <p>Antigen Retrieval: Boil tissue section in 10mM citrate buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min.</p> <p>Primary Antibody Incubation: Incubate for 30 minutes at RT.</p> <p>Slide Washing: Slides must be washed in between steps. Rinse slides with PBS/0.05% Tween.</p> <p>Visualization: Detect the antibody as instructed by the instructions provided with the visualization system.</p>
WESTERN BLOTTING:	Recommended starting protocol: Dilute the antibody 1:25. Incubate for 1 hour at room temperature. The dilution is an estimate; actual results may differ because of variability in methods and protocols. Optimal dilution and procedure should be determined by the end user.
POSITIVE CONTROL:	Esophageal carcinoma
CELLULAR LOCALIZATION:	Membrane
STORAGE & STABILITY	Store at 2-8°C. Do not freeze. The user must validate any other storage conditions. When properly stored, the reagent is stable to the date indicated on the label. Do not use the reagent beyond the expiration date. There are no definitive signs to indicate instability of this product; therefore, positive and negative controls should be tested simultaneously with unknown specimens.
WARNINGS & PRECAUTIONS:	<ol style="list-style-type: none"> Avoid contact of reagents with eyes and mucous membranes. If reagents come into contact with sensitive areas, wash with copious amounts of water. This product is harmful if swallowed. Consult local or state authorities with regard to recommended method of disposal. Avoid microbial contamination of reagents.



UK & Rest of World
184 Milton Park, Abingdon
OX14 4SE, Oxon, UK
Tel: +44 (0) 1235 828 200
Fax: +44 (0) 1235 820 482

Switzerland
Centro Nord-Sud 2E
CH-6934 Bioggio-Lugano
Tel: +41 (0) 91 604 55 22
Fax: +41 (0) 91 605 17 85

Deutschland
Bockenheimer Landstr. 17/19
60325 Frankfurt/Main
Tel: +49 (0) 69 779099
Fax: +49 (0) 69 13376880

United States
23591 El Toro Rd, Suite #167
Lake Forest, CA 92630
Tel: +1 800 987 0985
Fax: +1 949 265 7703