



Specification Sheet

Select-HA™ HiLadder

Catalogue No. HYA-HILAD-20

Product Description

Select-HA™ is a hyaluronic acid (HA) preparation of uniform and narrow size distribution prepared by *in vitro* synthesis using recombinant *Pasteurella multocida* hyaluronan synthase¹. The HiLadder contains five Select-HA™ molecular mass markers in the range of ~500 kDa to 1500kDa.

Quantity

The tube contains 25 µg (total) of five different Select-HA™ preparations supplied as a mixture of lyophilized sodium salts. The materials are sufficient for at least 20 lanes.

Recommended Reconstitution Procedure

Centrifuge the tube for a few seconds to collect the Select-HA™ solids in the bottom of the tube. Carefully open and add 100 µl of sterile water directly to the bottom of the tube. Allow two hours at 4°C for sample rehydration and then mix well before use. Electrophoresis of 5µl of reconstituted sample (Fig. 1) using a standard gel loading buffer on an agarose gel (0.6 – 1%) results in clearly defined bands when stained with 0.005% Stains-all (in 50% ethanol)².

Storage

Store at -20°C or below. To avoid frequent freeze-thaw, aliquoting is recommended. Avoid contamination with microbes or HA-degrading enzymes.

References

1. DeAngelis, P.L. *et al.* (1998) *J. Biol. Chem.*, **273**, 8454–8458.
2. Lee, H.G. *et al.* (1994) *Analytical Biochem.* **219**, 278-287.

For research use as molecular mass markers only

U.S. patent 6,444,447 and 7,223,571

AMSBIO | www.amsbio.com | info@amsbio.com



UK & Rest of the World
184 Park Drive, Milton Park
Abingdon OX14 4SE, UK
T: +44 (0)1235 828 200
F: +44 (0) 1235 820 482



North America
1035 Cambridge Street,
Cambridge, MA 02141
T: +1 (617) 945-5033 or
T: +1 (800) 987-0985
F: +1 (617) 945-8218



Germany
Bockenheimer Landstr. 17/19
60325 Frankfurt/Main
T: +49 (0) 69 779099
F: +49 (0) 69 13376880



Switzerland
Centro Nord-Sud 2E
CH-6934 Bioggio-Lugano
T: +41(0) 91 604 55 22
F: +41(0) 91 605 17 85