

Anti-ICAM2(CD102) aptamer, Magnetic Cell Isolation Kit (Catalog No. ICAM2-2296BCI/ ICAM2-2296FBCI)

Description

AptSci provides two types of kits. One is biotinylated aptamer based cell isolation kit (Catalog No. ICAM2-2296BCI). The other is bi-labeled (FITC dye conjugated biotin aptamer) aptamer based cell isolation kit (Catalog No. ICAM2-2296FBCI). Aptamer based magnetic cell isolation kit products do not adversely affect cells during isolation process, thus can be used to isolate pure, viable and functional cells which advance your biology research.

AptSci ICAM2 cell isolation kit is ideal for positive isolation of ICAM2 expressing target cells directly from all types of samples. Cell can also be eluted from bead-cell complexes with releasing buffer included in Kit, and then be used in all downstream experiments, including flow cytometry, cell culture and molecular studies.

Component description

- AptSci aptamer is a single stranded oligonucleotide that is engineered through advanced SELEX with modified nucleotide.
- Aptamer is generated with recombinant human ICAM2 protein produced in mammalian cells and binds their cellular target with high affinity and specificity (Fig. 1). ICAM2 aptamer cross-reacts with mouse ICAM2.
- Magnetic beads are uniform, colloiddally stable and non-porous beads (1µm diameter) covalently coupled with streptavidin.
- Bi-labeled aptamer that has FITC at 5'-end to monitor protein expression by flow cytometry and biotin at 3'-end to separate target cells.




Fig. 1. Flow cytometry histograms showing the binding of representative ICAM2 aptamer in Jurkat cells.

Approximately 1×10^6 cells were stained with FITC conjugated ICAM2 aptamer (Red histogram). As a control, the cells were stained FITC conjugated control aptamer (Gray histogram).

Principle of the AptoPrep™ Cell Isolation

AptSci cell isolation kit is designed to isolate cells via an indirect method and for positive selection principle using biotinylated aptamers and streptavidin magnetic beads.

Target cells are specifically labeled with biotinylated aptamer against cell surface target of desired cells. Streptavidin magnetic beads allow for efficient binding to the aptamer labeled cell. Magnetically labeled target cells are then separated from unlabeled cells using magnet. FACS analysis can immediately be performed with bi-labeled aptamer during cell isolation process (Catalog No. ICAM2-2296FBCI). In final step, bead-free and aptamer-free target cells were released from bead-bound cells (positive fraction) using releasing component (Fig. 2).

Positive isolation: Discard the supernatants and use the bead-bound cells for downstream application.

Release target cells from beads: Bead-bound cells are washed and target cells are released from the beads with releasing buffer included in Kit.





Fig. 2. Overview of AptoPrep cell isolation procedure.



UK & Rest of World
184 Milton Park, Abingdon
OX14 4SE, Oxon, UK
Tel: +44 (0) 1235 828 200
Fax: +44 (0) 1235 820 482

Switzerland
Centro Nord-Sud 2E
CH-6934 Bioggio-Lugano
Tel: +41 (0) 91 604 55 22
Fax: +41 (0) 91 605 17 85

Deutschland
Bockenheimer Landstr. 17/19
60325 Frankfurt/Main
Tel: +49 (0) 69 779099
Fax: +49 (0) 69 13376880

North America
23591 El Toro Rd, Suite #180
Lake Forest, CA 92630
Tel: +1 800 987 0985
Fax: +1 949 265 7703

amsbio
info@amsbio.com
www.amsbio.com
AMS Biotechnology

Anti-ICAM2(CD102) aptamer, Magnetic Cell Isolation Kit (Catalog No. ICAM2-2296BCI/ ICAM2-2296FBCI)

I. HUMAN APPLICATION

Typical results of the AptoPrep™ ICAM2 Cell Isolation

Isolation of ICAM2⁺ cells from fibroblast was performed with AptSci ICAM2 cell isolation kit. Human T lymphocyte Jurkat cells (ICAM2 positive cell, ~1x10⁶ cells) were spiked with MG-63 (ICAM2 negative cell, 1x10⁷ cells). Yield of ICAM2⁺ cells isolation was measured at 54%. Purity and viability of recovered ICAM2⁺ cells were measured at 96.4% and 83.9% (data not shown) respectively (Fig. 3).




Fig. 3. Isolation of ICAM2⁺ cells from ICAM2⁻ cells with AptSci ICAM2 cell isolation kit.

Both start sample before isolation and negative fraction after isolation were stained with FITC-ICAM2 aptamer (Gate of density plot). Positive fraction is stained with FITC-ICAM2 aptamer (Gate and red histogram). As a control, the cells were stained with FITC conjugated control aptamer.

Downstream application

ICAM2⁺ cells can be efficiently isolated from a sample with AptSci ICAM2 cell isolation kit. Lyse the cells directly after isolation, and isolate proteins, DNA, or mRNA to be used in PCR, microarrays, proteomics, and other applications where the removal of beads is not required. For functional studies such as cytokine expression, proliferation/apoptosis induction or for flow cytometry analysis, the cells need to be released from beads after positive isolation of cell. Releasing buffer included in Kit will allow you to collect the bead-free and aptamer-free ICAM2⁺ cell.

After elution of ICAM2⁺ cells with releasing buffer, elution yield was calculated by counting cells with hemocytometer and measured at 51% (data not shown). Purity and viability of released ICAM2⁺ cells were measured at 97.5% and 88.2%, respectively (Fig. 4).

Fig. 4. Flow cytometry showing bead-free and aptamer-free ICAM2⁺ cells are released with releasing buffer.

Released cell fraction (Blue histogram) was stained with FITC-ICAM2 aptamer and propidium iodide for cell viability (Red dot plot). As a control, the cells were stained with FITC conjugated control aptamer (Gray histogram).

II. MOUSE APPLICATION

Endothelial cell isolation from mouse lung tissue.

ICAM2 is a marker of endothelial cell. Isolation of endothelial cells from primary mouse lung cells were performed with AptSci ICAM2 cell isolation kit. Mouse primary cells were prepared from mouse lung tissue by disruption with collagenase. Yield of ICAM2⁺ cells isolation was measured at 52 % and purity of recovered ICAM2⁺ cells were measured at 89.1% (Fig. 5).

Fig. 5. Isolation of endothelial cell (ICAM2⁺ cells) from mouse lung tissue with AptSci ICAM2 cell isolation kit.

Both start sample before isolation and negative fraction after isolation were stained with FITC-ICAM2 aptamer (Gate of black dot plot). Positive fraction is stained with FITC-ICAM2 aptamer (Gate and blue histogram). As a control, the cells were stained with FITC conjugated control aptamer.

Anti-ICAM2(CD102) aptamer, Magnetic Cell Isolation Kit (Catalog No. ICAM2-2296BCI/ ICAM2-2296FBCI)

Product Information

- **Product name:** Anti-ICAM2 aptamer, Magnetic Cell Isolation Kit
- **Catalog number:** ICAM2-2296BCI (biotinylated aptamer based cell isolation kit), ICAM2-2296FBCI (bi-labeled aptamer based cell isolation kit)
- **Content:** ICAM2-2296BCI (Biotinylated aptamer, streptavidin-coated magnetic bead and buffer), ICAM2-2296FBCI (FITC dye conjugated biotin aptamer, streptavidin-coated magnetic bead and buffer)
- **Form:** Dried aptamer and bead in 50 mM Tris pH8.0, 150 mM NaCl, 0.05% NaN₃.
- **Protein source for generation of aptamer:** Recombinant protein produced in mammalian cells.
- **Specificity:** Anti-ICAM2 aptamer binds to human and mouse ICAM2. Cross reactivity with other species has not been tested.
- **MW:** ~26 kDa
- **Tested applications:** FACS and cell isolation
- **Shipping & Storage :** At 2°C to 8°C. There is no decrease in performance of the kit after storage for 1 year at 2°C to 8°C.

LIMITATIONS

Warranty: AptSci AptoPrep™ products are warranted to meet stated product specifications and to confirm to label descriptions when used and stored properly. Unless otherwise stated, this warranty is limited to one year from date of sales for products used, handled and stored according to AptSci's instructions. AptSci's sole liability is limited to replacement of the product or refund of the purchase price. AptoPrep™ products are supplied for research use only. They are not intended for medicinal, diagnostic or therapeutic use. AptoPrep™ may not be resold, modified for resale or used to manufacture commercial products without prior written approval from AptSci.



UK & Rest of World

184 Milton Park, Abingdon
OX14 4SE, Oxon, UK
Tel: +44 (0) 1235 828 200
Fax: +44 (0) 1235 820 482

Switzerland

Centro Nord-Sud 2E
CH-6934 Bioggio-Lugano
Tel: +41 (0) 91 604 55 22
Fax: +41 (0) 91 605 17 85

Deutschland

Bockenheimer Landstr. 17/19
60325 Frankfurt/Main
Tel: +49 (0) 69 779099
Fax: +49 (0) 69 13376880

North America

23591 El Toro Rd, Suite #180
Lake Forest, CA 92630
Tel: +1 800 987 0985
Fax: +1 949 265 7703

amsbio

info@amsbio.com

www.amsbio.com
AMS Biotechnology