

# Cathepsin S Activity Assay Kit

(Catalog #K144-100; 100 assays; Store kit at -20°C)

## I. Introduction:

Apoptosis can be mediated by mechanisms other than the traditional caspase-mediated cleavage cascade. There is growing recognition that alternative proteolytic enzymes such as the lysosomal cathepsin proteases may initiate or propagate proapoptotic signals. Cathepsins are lysosomal enzymes that are also used as sensitive markers in various toxicological investigations. The Cathepsin-S Activity Assay kit is a fluorescence-based assay that utilizes the preferred cathepsin-S substrate sequence VVR labeled with AFC (amino-4-trifluoromethyl coumarin). Cell lysates or other samples that contain cathepsin-S will cleave the synthetic substrate Z-VVR-AFC to release free AFC. The released AFC can easily be quantified using a fluorometer or fluorescence plate reader. The cathepsin-S assay is simple, straightforward, and can be adapted to 96-well plate assays. Assay conditions have been optimized to obtain the maximal activity.

## II. Kit Contents:

Components	K-140-100	Cap Code	Part No.
CS Cell Lysis Buffer	25 ml	WM	K144-100-1
CS Reaction Buffer	5 ml	NM	K144-100-2
CS Substrate Z-VVR-AFC (10 mM)	0.2 ml	Brown	K144-100-3
CS Inhibitor (1 mM)	20 µl	Red	K144-100-4

## III. Storage and Stability:

- Store kit at -20°C (Store CS Cell Lysis Buffer and CS Reaction Buffer at 4°C after opening). Protect CS Substrate Z-VVR-AFC from light. All reagents are stable for 6 months under proper storage conditions.

## IV. Cathepsin S Assay Protocol:

- Collect cells ( $1-5 \times 10^6$ ) by centrifugation.  
Note: Use 50-200 µg cell lysates (in 50 µl of CS Cell Lysis Buffer) if protein concentration has been measured.
- Lyse cells in 50 µl of chilled CS Cell Lysis Buffer. Incubate cells on ice for 10 min.
- Centrifuge at top speed in a microcentrifuge for 5 min, transfer the supernatant to a new tube. Add 50 µl of cell lysate to a 96-well plate.
- Add 50 µl of CS Reaction Buffer to each sample.
- Add 2 µl of the 10 mM Z-VVR-AFC substrate (200 µM final concentration).  
Note: For negative control, add 2 µl of CS Inhibitor prior to adding CS Substrate, or make a reaction mixture that does not contain sample as control.
- Incubate at 37°C for 1-2 hour.
- Read samples in a fluorometer equipped with a 400-nm excitation filter and 505-nm emission filter. For a plate-reading set-up, transfer the samples to a 96-well plate. You may also perform the entire assay directly in a 96-well plate.

Fold-increase in Cathepsin S activity can be determined by comparing the relative fluorescence units (RFU) with the level of the uninduced control or the negative control sample. If desired, the units of cathepsin S can be determined by generating a standard curve using free AFC under your assay conditions. Free AFC is available from BioVision (Cat.# 1077-100).

## V. Related Products:

### Apoptosis Detection Kits & Reagents

- Annexin V Kits & Bulk Reagents
- Caspase Assay Kits & Reagents
- Mitochondrial Apoptosis Kits & Reagents
- Nuclear Apoptosis Kits & Reagents
- Additional Apoptosis Kits & Reagents

### Cell Fractionation System

- Mitochondria/Cytosol Fractionation Kit
- Nuclear/Cytosol Fractionation Kit
- Membrane Protein Extraction Kit
- Cytosol/Particulate Rapid Separation Kit
- Mammalian Cell Extraction Kit
- FractionPREP Fractionation System

### Cell Proliferation & Senescence

- Quick Cell Proliferation Assay Kit
- Senescence Detection Kit
- High Throughput Apoptosis/Cell Viability Assay Kits
- LDH-Cytotoxicity Assay Kit
- Bioluminescence Cytotoxicity Assay Kit
- Live/Dead Cell Staining Kit

### Cell Damage & Repair

- HDAC Fluorometric & Colorimetric Assays & Drug Discovery Kits
- HAT Colorimetric Assay Kit & Reagents
- DNA Damage Quantification Kit
- Glutathione Fluorometric & Colorimetric Assay Kits
- Nitric Oxide Fluorometric & Colorimetric Assay Kits

### Signal Transduction


- cAMP & cGMP Assay Kits
- Akt & JNK Activity Assay Kits
- Beta-Secretase Activity Assay Kit

### Adipocyte & Lipid Transfer

- Recombinant Adiponectin, Survivin, & Leptin
- CETP and PLTP Activity Assay & Drug Discovery Kits
- Total Cholesterol Quantification Kit

### Molecular Biology & Reporter Assays

- siRNA Vectors
- Cloning Insert Quick Screening Kit
- Mitochondrial & Genomic DNA Isolation Kits
- 5 Minutes DNA Ligation Kit
- 20 Minutes Gel Staining/Destaining Kit



#### UK & Rest of World

184 Milton Park, Abingdon  
OX14 4SE, Oxon, UK  
Tel: +44 (0) 1235 828 200  
Fax: +44 (0) 1235 820 482

#### Switzerland

Centro Nord-Sud 2E  
CH-6934 Bioggio-Lugano  
Tel: +41 (0) 91 604 55 22  
Fax: +41 (0) 91 605 17 85

#### Deutschland

Bockenheimer Landstr. 17/19  
60325 Frankfurt/Main  
Tel: +49 (0) 69 779099  
Fax: +49 (0) 69 13376880

#### North America

23591 El Toro Rd, Suite #167  
Lake Forest, CA 92630  
Tel: +1 800 987 0985  
Fax: +1 949 265 7703


amsbio

info@amsbio.com

www.amsbio.com  
AMS BioTechnology

## GENERAL TROUBLESHOOTING GUIDE

Problems	Cause	Solution
Assay not working	<ul style="list-style-type: none"> <li>Cells did not lyse completely</li> <li>Experiment was not performed at optimal time after apoptosis induction</li> <li>Plate read at incorrect wavelength</li> </ul>	<ul style="list-style-type: none"> <li>Resuspend the cell pellet in the lysis buffer and incubate as described in the datasheet</li> <li>Perform a time-course induction experiment for apoptosis</li> <li>Check the wavelength listed in the datasheet and the filter settings of the instrument</li> </ul>
High Background	<ul style="list-style-type: none"> <li>Increased amount of cell lysate used</li> <li>Increased amounts of components added due to incorrect pipetting</li> <li>Incubation of cell samples for extended periods</li> <li>Use of expired kit or improperly stored reagents</li> <li>Contaminated cells</li> </ul>	<ul style="list-style-type: none"> <li>Refer to datasheet and use the suggested cell number to prepare lysates</li> <li>Use calibrated pipettes</li> <li>Refer to datasheet and incubate for exact times</li> <li>Always check the expiry date and store the individual components appropriately</li> <li>Check for bacteria/ yeast/ mycoplasma contamination</li> </ul>
Lower signal levels	<ul style="list-style-type: none"> <li>Cells did not initiate apoptosis</li> <li>Very few cells used for analysis</li> <li>Use of samples stored for a long time</li> <li>Incorrect setting of the equipment used to read samples</li> <li>Allowing the reagents to sit for extended times on ice</li> </ul>	<ul style="list-style-type: none"> <li>Determine the time-point for initiation of apoptosis after induction (time-course experiment)</li> <li>Refer to datasheet for appropriate cell number</li> <li>Use fresh samples or aliquot and store and use within one month for the assay</li> <li>Refer to datasheet and use the recommended filter setting</li> <li>Always thaw and prepare fresh reaction mix before use</li> </ul>
Samples with erratic readings	<ul style="list-style-type: none"> <li>Uneven number of cells seeded in the wells</li> <li>Samples prepared in a different buffer</li> <li>Adherent cells dislodged and lost at the time of experiment</li> <li>Cell/ tissue samples were not completely homogenized</li> <li>Samples used after multiple freeze-thaw cycles</li> <li>Presence of interfering substance in the sample</li> <li>Use of old or inappropriately stored samples</li> </ul>	<ul style="list-style-type: none"> <li>Seed only equal number of healthy cells (correct passage number)</li> <li>Use the cell lysis buffer provided in the kit</li> <li>Perform experiment gently and in duplicates/triplicates; apoptotic cells may become floaters</li> <li>Use Dounce homogenizer (increase the number of strokes); observe efficiency of lysis under microscope</li> <li>Aliquot and freeze samples, if needed to use multiple times</li> <li>Troubleshoot as needed</li> <li>Use fresh samples or store at correct temperatures until use</li> </ul>
Unanticipated results	<ul style="list-style-type: none"> <li>Measured at incorrect wavelength</li> <li>Cell samples contain interfering substances</li> </ul>	<ul style="list-style-type: none"> <li>Check the equipment and the filter setting</li> <li>Troubleshoot if it interferes with the kit (run proper controls)</li> </ul>
General issues	<ul style="list-style-type: none"> <li>Improperly thawed components</li> <li>Incorrect incubation times or temperatures</li> <li>Incorrect volumes used</li> <li>Air bubbles formed in the well/tube</li> <li>Substituting reagents from older kits/ lots</li> <li>Use of a different 96-well plate</li> </ul>	<ul style="list-style-type: none"> <li>Thaw all components completely and mix gently before use</li> <li>Refer to datasheet &amp; verify the correct incubation times and temperatures</li> <li>Use calibrated pipettes and aliquot correctly</li> <li>Pipette gently against the wall of the well/tubes</li> <li>Use fresh components from the same kit</li> <li>Fluorescence: Black plates; Absorbance: Clear plates</li> </ul>
<b>Note#</b> The most probable cause is listed under each section. Causes may overlap with other sections.		



### UK & Rest of World

184 Milton Park, Abingdon  
OX14 4SE, Oxon, UK  
Tel: +44 (0) 1235 828 200  
Fax: +44 (0) 1235 820 482

### Switzerland

Centro Nord-Sud 2E  
CH-6934 Bioggio-Lugano  
Tel: +41 (0) 91 604 55 22  
Fax: +41 (0) 91 605 17 85

### Deutschland

Bockenheimer Landstr. 17/19  
60325 Frankfurt/Main  
Tel: +49 (0) 69 779099  
Fax: +49 (0) 69 13376880

### North America

23591 El Toro Rd, Suite #167  
Lake Forest, CA 92630  
Tel: +1 800 987 0985  
Fax: +1 949 265 7703