


Rabbit Anti-Human CD79a Monoclonal Antibody (Clone SP18)

CATALOG #:

M3180 0.1 ml rabbit monoclonal antibody supplied as tissue culture supernatant in TBS/1% BSA buffer pH 7.5 with less than 0.1% sodium azide.

M3184 1 ml rabbit monoclonal antibody supplied as tissue culture supernatant in TBS/1% BSA buffer pH 7.5 with less than 0.1% sodium azide.

M3181 7 ml pre-diluted rabbit monoclonal antibody supplied as culture supernatant in TBS/1% BSA buffer pH 7.6 with less than 0.1% sodium azide.



Human B cell lymphoma stained with anti-CD79a antibody

INTENDED USE: For Research Use Only. Not for use in diagnostic procedures.

CLONE: SP18

IMMUNOGEN: Synthetic peptide derived from N-terminus region of human CD79a protein.

IG ISOTYPE: Rabbit IgG

EPITOPE: Not determined

MOLECULAR WEIGHT 50kDa

SPECIES REACTIVITY: Human (tested). Monkey, horse, cow, pig, rabbit, guinea pig, rat and chicken (by sequence homology). It does not react with opossum. Others not tested.

DESCRIPTION: A disulphide-linked heterodimer, consisting of CD79a / mb-1 and CD79b / B29 polypeptides, CD79a is non-covalently associated with membrane-bound immunoglobulins on B cells to constitute the B cell Ag receptor. CD79a first appears at pre B cell stage and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas.

APPLICATIONS: Immunohistochemistry (IHC)

IHC PROCEDURE:

Specimen Preparation: Formalin-fixed, paraffin-embedded tissues are suitable for use with this primary antibody.

Deparaffinization: Deparaffinize slides using xylene or xylene alternative and graded alcohols.

Antibody Dilution: If using the concentrate format of this product, dilute the antibody 1:200. The dilutions are estimates; actual results may differ because of variability in methods and protocols.

Antigen Retrieval: Boil tissue section in 10mM citrate buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min.

Primary Antibody Incubation: Incubate for 30 minutes at RT.

Slide Washing: Slides must be washed in between steps. Rinse slides with PBS/0.05% Tween.

Visualization: Detect the antibody as instructed by the instructions provided with the visualization system.

POSITIVE CONTROL: Tonsil and B cell lymphoma

CELLULAR LOCALIZATION: Membrane and cytoplasm

STORAGE & STABILITY Store at 2-8°C. Do not freeze. The user must validate any other storage conditions. When properly stored, the reagent is stable to the date indicated on the label. Do not use the reagent beyond the expiration date.

There are no definitive signs to indicate instability of this product; therefore, positive and negative controls should be tested simultaneously with unknown specimens.

WARNINGS & PRECAUTIONS:

1. Avoid contact of reagents with eyes and mucous membranes. If reagents come into contact with sensitive areas, wash with copious amounts of water.
2. This product is harmful if swallowed.
3. Consult local or state authorities with regard to recommended method of disposal.
4. Avoid microbial contamination of reagents.

