

PRODUCT INFORMATION

Collagenase AF-1 GMP Grade (≥ 150 U)

Cat. No. N0003855

Introduction Collagenases from *Clostridium histolyticum* are proteolytic enzymes that cleave peptide bonds in the triple helical collagen molecules of human or animal tissue *in situ*.

For this reason collagenases are widely used for isolation of various cell types by tissue dissociation.

Description Collagenase AF-1 GMP Grade is chromatographically highly purified leading to a very high collagenolytic activity. It is largely free from additional enzymatic activities like clostripain, trypsin-like activities and neutral protease, as well as endotoxins.

Collagenase AF-1 GMP Grade is manufactured according to GMP guidelines using a production process completely free of animal-based components. Thus, the introduction of any potential animal-derived pathogen is excluded by design.

The absence of any animal-derived ingredients and thorough microbial analysis provide the highest possible safety.

Specification

Collagenase activity	≥ 3.000 U/mg (PZ acc. to Wunsch)
	≥ 150 U/vial (PZ acc. to Wunsch)
Neutral protease activity	≤ 0.050 U/mg (DMC)
Trypsin-like activities	status
Clostripain, native state	status
TAMC	≤ 10 /vial
TYMC	≤ 10 /vial
Bacterial endotoxins	≤ 10.0 EU/mg

Application Collagenase AF-1 GMP Grade is suitable for cell isolation from several tissue types intended for clinical applications.

For larger requirements, Collagenase AF-1 GMP Grade is also available in a pack size of ≥ 2000 PZ U per vial (Cat. No. N0003554).

Collagenase AF-1 GMP Grade is often used in combination with Neutral Protease AF GMP Grade (Cat. No. N0003553).

Storage conditions Collagenase AF-1 GMP Grade is available as a lyophilized powder. It should be stored at +2 to +8 °C in a dry environment.

Under these conditions the product is stable until the date stated on the certificate of analysis if repeated opening and closing of the vial is avoided.

Documents A certificate of analysis is provided for each lot.

For additional documentation (e.g. stability data, GMP certificate, certificate of origin, etc.), please contact our product management team at collagenase@nordmark-biochemicals.com

Product size

Product	Cat. No.	Size (PZ U)
Collagenase AF-1 GMP Grade	N0003855	≥ 150
	N0003554	≥ 2000

www.amsbio.com | info@amsbio.com

PRODUCT INFORMATION

Collagenase AF-1 GMP Grade (≥ 150 U)

Cat. No. N0003855

Instructions for use:

General Collagenase AF-1 GMP Grade, often in combination with Neutral Protease AF GMP Grade (Cat. No. N0003553), is suitable for the dissociation of sensitive cells from several tissues, such as liver, pancreas and tumor. In addition, it has been shown to be highly effective for the isolation of the stromal vascular fraction (SVF) from adipose tissue.

Tissue dissociation Recommended final concentrations for isolation of SFV from human lipoaspirate:

Collagenase AF-1 GMP Grade: 0.2 PZ U/ml
 Neutral Protease AF GMP Grade: 0.1 DMC U/ml

Collagenase AF-1 GMP Grade is provided in vials with ≥ 150 PZ U which is usually sufficient for approx. 800 ml lipoaspirate.

The appropriate collagenase concentration depends on the tissue type and origin as well as on the applied isolation procedure.

Collagenase activity is at an optimum at 37 °C and pH 7.4.

Stock solution A stock solution of Collagenase AF-1 GMP Grade can be prepared by dissolving the enzyme in buffer. The enzyme solution should be constantly stored on ice.

Since collagenase depends on calcium, absolutely no calcium chelating agents (e.g. EDTA) should be present at all. A buffer with ≥ 2 mM Ca^{2+} is recommended to be used.

Reconstituted Collagenase AF-1 GMP Grade can be filtered, aliquoted and stored at -20 °C for up to 1 year. Repeated freezing and thawing should be avoided. For 0.22 μm filtration filters with low protein binding properties (e.g. cellulose acetate, PVDF, or PES) are recommended.

Working solution To prepare a working solution, the stock solution is diluted to achieve the desired collagenase concentration.

The working solution should be constantly stored on ice until use.

If Collagenase AF-1 GMP Grade solution is mixed with neutral protease solution, the blend should be used immediately.

Inactivation and inhibitors The dissociation process can be reduced, e.g. by cooling down or diluting the enzyme solution.
 Collagenase is reversibly inactivated at high pH values and irreversibly inactivated at low pH values. Inhibitors of collagenase include cysteine or chelating agents like EDTA.

Important note Collagenase AF-1 GMP Grade is not intended for direct application in humans. It is suitable for cell isolation from several tissue types intended for clinical application or transplantation into humans.

AMS BIO | www.amsbio.com | info@amsbio.com