

GenePORTER[®] 2

amsbio

Transfection Reagent

Catalog #	Contents	Quantity
T202007 (75 reactions)	GenePORTER [®] 2 Lipid Film, dried	1 vial
	Hydration Buffer, Transfection Grade	1 x 0.8 ml
	DNA Diluent	1 x 4 ml
	DNA Diluent B	1 x 4 ml
T202015 (150 reactions)	GenePORTER [®] 2 Lipid Film, dried	1 vial
	Hydration Buffer, Transfection Grade	1 x 1.6 ml
	DNA Diluent	1 x 8 ml
	DNA Diluent B	1 x 8 ml
T202075 (750 reactions)	GenePORTER [®] 2 Lipid Film, dried	5 vials
	Hydration Buffer, Transfection Grade	5 x 1.6 ml
	DNA Diluent	5 x 8 ml
	DNA Diluent B	5 x 8 ml
Shipping	Shipped at room temperature.	
Storage	Store at 4°C; stable for 1 year.	

Related Products	Catalog #
GenePORTER [®] 2 Transfection Reagent, 150 reactions	T202015
GenePORTER [®] 2 Transfection Reagent, 750 reactions	T202075
GenePORTER [®] Transfection Reagent, 75 reactions	T201007
GenePORTER [®] Transfection Reagent, 150 reactions	T201015
GenePORTER [®] Transfection Reagent, 750 reactions	T201075
BoosterExpress [™] Reagents, 3 x 1.5 ml	T20100B
GeneSilencer [®] siRNA Transfection Reagent, 200 reactions	T500750
GeneSilencer [®] siRNA Transfection Reagent, 1000 reactions	T505750
NeuroFect [™] Transfection Reagent, 75-300 reactions	T800075
NeuroFect [™] Transfection Reagent, 375-1500 reactions	T800750
BioPORTER [®] Protein Delivery Reagent, 24 reactions	BP502401
BioPORTER [®] Protein Delivery Reagent, QuikEase kit, 24 rxns.	BP502424
Cytofectin [™] Transfection Reagent, 1 ml	T610001
Cytofectin [™] Transfection Reagent, 5 x 1 ml	T610005

Introduction: The GenePORTER[®] 2 transfection reagent is the most effective and widespread gene delivery reagent developed by Genlantis. While offering all of the advantages of DHC technology as the original GenePORTER[®] Reagent, the GenePORTER 2 Reagent delivers higher transfection efficiencies for more difficult to transfect cell types. With a choice of two optimized DNA diluent buffers, GenePORTER 2 performs effectively across a broad range of cells, offers the highest efficiencies, and is effective in the presence of serum, eliminating inconvenient media changes.

METHODS AND PROCEDURES

A. DNA DILUENT SELECTION

Select the DNA diluent suitable for your cell type as listed below; if not listed, choose the cell type that closely matches yours or contact us for a citation list at info@amsbio.com. If no data is available for your cell type, try the DNA Diluent B protocol first (Section C).

Cell Lines	DNA Diluent	DNA Diluent B	Serum
HeLa-S3	★	★★	○
HeLa	★	★★	○
COS-1	★	★	●
COS-7	★	★	●
Hep-G2	★	★	●
NIH-3T3	★	★	●
MDCK	★	★★	○
K-562	★	★★	○
CV-1	★	★	●
B15-F0	★	★	●
293	★	★	●
BHK-21	★	★	●
CHO-K1	★▲	★▲	●
PC-12	★	NR	●
P19	★	★	●
HUVEC-C	★	★	●
Jurkat	◆	◆	○

Legend

★ Works well	○ Works well w/o serum
★★ Works better	● Works well w/ and w/o serum
NR Not recommended	
▲ Best expression is w/o serum during 1 st hr. of transfection	
◆ Original GenePORTER [®] reagent is recommended	

B. GENEPORTER LIPID HYDRATION

- Hydrate each GenePORTER 2 lipid vial at room temperature with 0.75 ml (for T202007) or 1.5 ml (for T202015 and T202075) of the Hydration Buffer. Vortex for 10 seconds at top speed before use. Store hydrated reagent at 4°C; vortex briefly before each use.
- Use 25 µl of the DNA Diluent or DNA Diluent B per each 1 µg of DNA. Avoid vortexing either of the DNA diluent solutions.

C. PROTOCOL FOR USING THE DNA DILUENT

- For most cell types, use 5 µl GenePORTER 2 per 1 µg of DNA.
- Your DNA can be suspended in TE buffer or purified water. A DNA concentration of at least 0.1 mg/ml works well for most reaction sizes.

i. Transfection of Adherent Cells

- Dilute the hydrated GenePORTER 2 reagent with serum-free medium as shown in Table 1 below.
- Dilute the DNA with DNA Diluent as shown in Table 1 below; mix well by pipetting and incubate at room temperature for 1-5 minutes.
NOTE: Do not vortex DNA Diluent.

Table 1: Amounts of DNA, diluent, GenePORTER 2, and Medium

DNA (µg)	DNA Diluent (µl)	GenePORTER 2 (µl)	Serum-free Medium (µl)
0.5	12.5	2.5	10.0
1.0	25.0	5.0	20.0
2.0	50.0	10.0	40.0
4.0	100.0	20.0	80.0
8.0	200.0	40.0	160.0

- Add the diluted DNA to the diluted GenePORTER 2 reagent. Incubate at room temperature for 5 to 10 minutes to form lipid/DNA complexes (lipoplexes).
NOTE: Do not incubate for more than 30 minutes.

8. Add the GenePORTER 2/DNA complexes directly to cells growing in serum-containing medium and incubate at 37°C. Use Table 2 below for recommended transfection volumes and DNA amounts.

NOTES: Cells plated the day before transfection should be 50%-70% confluent on transfection day. Omitting antibiotics from the media during transfection may increase expression levels; this effect is cell-type dependent and usually small.

For some cells (such as HeLa S3, MDCK, CHO-K1), higher transfection efficiencies can be achieved when the initial 4-hour incubation is done in serum-free media, followed by the addition of one volume of medium containing 20% serum.

Table 2: Transfection Volumes and DNA Amounts Per Dish Size

Tissue Culture Dish Size	DNA (µg)	Transfection Volume (ml)
96-well	0.1-0.5	0.1
24-well	0.5-2.0	0.25
6-well	2.0-6.0	1.0
60 mm	6.0-8.0	2.5
100 mm	8.0-12.0	5.0

9. Add fresh growth media as needed 24 hours post transfection; for some cell types, the old media can be replaced with fresh media at this step.
10. Depending on cell type and promoter activity, reporter gene assay can be performed 24-72 hours post transfection.
NOTE: the same protocol can be used to produce stably transfected cells: 48 to 72 hours post transfection, put the cells in fresh medium containing the appropriate selection antibiotic. Wait at least 48 hours before exposing the transfected cells to the selection medium, and for some cell types it may be necessary to wait as long as 4 to 5 days.

ii. Transfection of Suspension Cells

GenePORTER 2 reagent works well for most suspension cells, such as K562 and PC 12, but for Jurkat cells we recommend using the original GenePORTER® Reagent.

For suspension cells, the protocol is the same as described for adherent cells, with the following exceptions:

11. The day before transfection, split the cells so they are in good condition on the day of transfection.
12. While the GenePORTER 2/DNA complexes are incubating, spin down the cells, resuspend them at 1×10^6 or 2×10^6 cells/ml in medium with or without serum, and transfer the appropriate volume to the dish as indicate in Table 3 below.
13. Add the GenePORTER 2/DNA complexes directly to the cells, and mix well by gently pipetting 2 to 3 times.
NOTE: this step is important because some suspension cells have a tendency to clump, which reduces transfection efficiency.
14. Incubate at 37°C and proceed as described for adherent cells.
NOTE: for some hematopoietic cell lines, mitogenic agents like PHA or PMA may be added to the cells 4 hours after transfection to a final concentration of 1 µg/ml or 50 ng/ml, respectively, to enhance the levels of gene expression.

Table 3: Number of Cells and Transfection Volumes for Suspension Cells

Tissue Culture Dish Size	Number of Cells (µg)	Transfection Volume (ml)
96-well	1×10^5	0.1
24-well	0.5×10^6	0.25
6-well	2×10^6	1.0
60 mm	5×10^6	2.5
100 mm	1×10^7	5.0

D. PROTOCOL FOR USING THE DNA DILUENT B

15. For most cell types, use 3.5 µl of the GenePORTER 2 reagent per 1 µg of DNA.

iii. Transfection of Adherent Cells

16. Dilute the hydrated GenePORTER 2 reagent with serum-free medium as shown in Table 4 below.
17. Dilute the DNA with DNA Diluent B as shown in Table 4 below; mix well by pipetting and incubate at room temperature for 5 minutes.
NOTE: Do not vortex DNA Diluent B.

Table 4: Amounts of DNA, diluent, GenePORTER 2, and Medium

DNA (µg)	DNA Diluent B (µl)	GenePORTER 2 (µl)	Serum-free Medium (µl)
0.5	12.5	1.75	10.75
1.0	25.0	3.5	21.5
2.0	50.0	7.0	43.0
4.0	100.0	14.0	86.0
8.0	200.0	28.0	172.0

18. Add the diluted DNA to the diluted GenePORTER 2. Incubate at room temperature for 5 minutes to form lipid/DNA complexes (lipoplexes).
NOTE: Do not incubate for more than 30 minutes.
19. Same as Step 8 above.
20. Same as Step 9 above.
21. Same as Step 10 above.

iv. Transfection of Suspension Cells

Same as Section ii protocol for transfection of suspension cells when using the DNA Diluent.

E. DETECTION OF EXPRESSED REPORTER GENES

β-Galactosidase

The following protocol¹ is provided for your convenience (alternatively, you could use one of the β-galactosidase assay kits offered by Genlantis Catalog Numbers: A10100K, A10200K, and A10300K):

Briefly, aspirate the culture media post transfection. Lyse the transfected cells from each well of a 96-well plate with 50 µl of the lysis buffer [0.1% Triton X-100 (w/v) in 250 mM Tris-HCl, pH 8.0], then subject the cells to one freeze-thaw cycle (freeze at -70°C and thaw at room temperature). While the cells are being lysed, prepare a β-galactosidase (*E.coli*; Sigma) standard curve with 0.5% BSA in PBS (w/v). Once the plate of lysed cells is completely thawed, transfer a 50-µl aliquot of each point on the standard curve to control wells of the plate. Typically, β-galactosidase expression ranges from 10,000 to 2,000,000 pg. Develop color by adding 150 µl of 1 mg/ml chlorophenol red-β-D-galactopyranoside (CPRG; Boehringer Mannheim) dissolved in β-gal buffer (1 mM MgCl₂; 10 mM KCl; 50 mM β-mercaptoethanol; and 60 mM Na₂HPO₄, pH 8.0). Allow the reaction to proceed at room temperature until the red color develops (2 min to 4 hours, depending on cell type). Read absorbance at 580 nm.

An immunohistochemical approach for quantifying β-galactosidase has also been reported³.

Green Fluorescent Protein

When green fluorescent protein (GFP) is the reporter gene used for transfections, use epifluorescence or confocal microscopy to detect expression. GFP has an excitation peak at 470 to 490 nm and emission peak at 510 nm. Expression levels of GFP can also be monitored by fluorescence-activated cell sorter analysis (FACS)⁴.

Secreted Alkaline Phosphatase

When heat-stable secreted alkaline phosphatase (SEAP) is the reporter gene used for transfections, use the following assay: heat supernatants from transfected cells at 65°C for 30 min to inactivate endogenous alkaline phosphatase activity. The SEAP transgene is stable during this treatment. Take aliquots of the culture media 48 hours posttransfection, and determine the SEAP activity quantitatively by using a colorimetric assay based on hydrolysis of the chromogenic substrate para-nitrophenyl phosphate (PNPP). Dissolve 1 mg/ml of PNPP reagent in a solution of 1 mM MgCl₂ and 100 mM diethanolamine, pH 9.8. Add 10 µl of 0.05% Zwittergent in PBS (free Ca²⁺ and Mg²⁺) into each well of a 96-well plate. Then add 20 µl of the heated cell culture media to each well. For control wells, 20 µl of water is used to normalize the volume. An alkaline phosphatase standard (EIA grade calf intestine alkaline phosphatase; Boehringer Mannheim) can be used to generate a standard curve from 10 to 10,000 pg per well. Add 200 µl of the PNPP substrate to each well to start the enzymatic reaction. Allow the reaction to incubate at room temperature until the yellow color develops. Using 0.05% Zwittergent in PBS as the diluent virtually reduces the background to zero, which increases the detection limit of the assay. Read the plates at 405 nm using either kinetic or static mode.

F. OPTIONAL PROTOCOL FOR LOW QUANTITY DNA TRANSFECTION

The following revised protocol^{1,2} can be used to facilitate pipetting and transfer of DNA/lipids complexes to the cells when a low quantity of DNA ($\leq 1 \mu\text{g}$) is used for transfection.

1. Dilute hydrated GenePORTER 2 reagent with serum-free medium as in Table 5 below.
2. First dilute the DNA diluent in serum-free medium and then add the DNA. See Table 5 (A, B, and C) below for volumes of serum-free medium, DNA diluent, and DNA amounts.

Table 5: Recommended Amounts of Reagents for Optional Protocol

A: Dilution of GenePORTER 2 Reagent		
DNA (μg)	Serum-free Medium (μl)	GenePORTER 2 Reagent (μl)
0.125	49.37	0.63
0.25	48.75	1.25
0.5	47.5	2.5
1.0	45.0	5.0

B. DNA Dilution		
Serum-free Medium (μg)	DNA Diluent (μl)	DNA (μg)
46.8	3.12	0.125
43.75	6.25	0.25
37.5	12.5	0.5
25.0	25.0	1.0

C. Transfection Volume and DNA Amounts Per Dish Size		
Tissue Culture Dish	DNA (μg)	Transfection Volume (ml)
96-well	0.1-0.25	0.1
24-well	0.5-2.0	0.25


3. Incubate 1 to 5 minutes at room temperature.
4. Proceed as in Steps 6 through 9 under the "Transfection of Adherent Cells" Section (when using the DNA Diluent).

REFERENCES

1. Felgner, JH. *et al.* (1994) Enhanced gene delivery and mechanism studies with a novel series of cationic lipid formulations. *J Biol Chem* **269**: 2550-2561.
2. Felgner, PL. *et al.* (1987) Lipofection: a highly efficient, lipid-mediated DNA-transfection procedure. *Proc Natl Acad Sci USA* **84**: 7413-7417.
3. Gussoni, E. *et al.* (1996) A method to codetect introduced genes and their products in gene therapy protocols. *Nature Biotechnology* **14**: 1012-1015.
4. Cheng, L. *et al.* (1996) Use of green fluorescent protein variants to monitor gene transfer and expression in mammalian cells. *Nature Biotechnology* **14**: 606-609.

LIMITED LICENSE: The purchase price paid for the GenePORTER® 2 Transfection Reagent Kits (hereto "GenePORTER 2") grants end users a non-transferable, non-exclusive license to use the kits and/or their components for **internal research use only** as described in this manual; in particular, research use only excludes and without limitation, resale, repackaging, or use for the making or selling of any commercial product or service without the written approval of Genlantis, a division of Gene Therapy Systems, Inc. (GTS) -- separate licenses are available for non-research use or applications. GenePORTER 2 and/or its components are not to be used for human diagnostic or included/used in any drug intended for human use. Care and attention should be exercised in handling the kit components by following appropriate research laboratory practices.

Purchasers may refuse this license by returning the enclosed materials unused. By keeping or using the enclosed materials, you agree to be bound by the terms of this license. The laws of the State of California shall govern the interpretation and enforcement of the terms of this License.



UK & Rest of World

184 Milton Park, Abingdon
OX14 4SE, Oxon, UK
Tel: +44 (0) 1235 828 200
Fax: +44 (0) 1235 820 482

Switzerland

Centro Nord-Sud 2E
CH-6934 Bioggio-Lugano
Tel: +41 (0) 91 604 55 22
Fax: +41 (0) 91 605 17 85

Deutschland

Bockenheimer Landstr. 17/19
60325 Frankfurt/Main
Tel: +49 (0) 69 779099
Fax: +49 (0) 69 13376880

United States

23591 El Toro Rd, Suite #167
Lake Forest, CA 92630
Tel: +1 800 987 0985
Fax: +1 949 265 7703

amsbio
info@amsbio.com