

Name: Purified Beta-catenin mouse monoclonal antibody, clone 2H3**Catalog:** TA500049**Product Data Sheet**

Gene Name: Homo sapiens catenin (cadherin-associated protein), beta 1, 88kDa (CTNNB1), transcript variant 1
GeneBank accession: NM_001904
IsoType: IgG2a

Reactivity: Human, Monkey
Test application: WB
Clone Name: clone 2H3

Gene Synonym: CTNNB; DKFZp686D02253; FLJ25606; FLJ37923

Validation Data:**Western Blot**

Guaranteed Applications: WB

Suggested Dilutions: Western blot: 1:1000

Immunogen: Recombinant protein expressed in E.coli corresponding to amino acids 531-781 of human beta-catenin

Components:

- Purified Beta-catenin mouse monoclonal antibody, clone 2H3 (TA500049)
- 1 vial of 20ug myc-DDK tagged CTNNB1 HEK293T over-expression lysate lyophilized in RIPA buffer (LY419662) (Reconstitute into 20ul of 1x SDS sample buffer before loading) (Not available for 30ul sample size)

Amount:

TA500049-100 100 uL

Concentration: 0.7mg/ml

Storage Condition: Shipped at 4C. Upon delivery store at -20C. Dilute in PBS (pH7.3) before use. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.


Buffer: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Purification:

Purified from mouse ascites fluids by affinity chromatography

Background:

Beta-catenin is an adherens junction protein. Adherens junctions (AJs; also called the zonula adherens) are critical for the establishment and maintenance of epithelial layers, such as those lining organ surfaces. AJs mediate adhesion between cells, communicate a signal that neighboring cells are present, and anchor the actin cytoskeleton. In serving these roles, AJs regulate normal cell growth and behavior. At several stages of embryogenesis, wound healing, and tumor cell metastasis, cells form and leave epithelia. This process, which involves the disruption and reestablishment of epithelial cell-cell contacts, may be regulated by the disassembly and assembly of AJs. AJs may also function in the transmission of the 'contact inhibition' signal, which instructs cells to stop dividing once an epithelial sheet is complete



Beta-catenin antibody (2H3) at 1:1000 dilution + 293T cell lysate

Related Product:

TrueORF cDNA clones
 VERIFY Tagged Antigen lysates
 HuSH-29 shRNA
 Western Blot reagents
 Anti-myc/DDK tag antibodies

This product is to be used for laboratory only. Not for diagnostic or therapeutic use.

UK & Rest of World

184 Milton Park, Abingdon
 OX14 4SE, Oxon, UK
 Tel: +44 (0) 1235 828 200
 Fax: +44 (0) 1235 820 482

Switzerland

Centro Nord-Sud 2E
 CH-6934 Bioggio-Lugano
 Tel: +41 (0) 91 604 55 22
 Fax: +41 (0) 91 605 17 85

Deutschland

Bockenheimer Landstr. 17/19
 60325 Frankfurt/Main
 Tel: +49 (0) 69 779099
 Fax: +49 (0) 69 13376880

United States

23591 El Toro Rd, Suite #167
 Lake Forest, CA 92630
 Tel: + 1 800 987 0985
 Fax: + 1 949 265 7703